


Earthworm Identification Guide

This guide can be used for the OPAL Soil and Earthworm Survey



Start here
Note that line drawings show the typical sizes of adult earthworms



Is it more than 2cm long, **AND** does it have a clearly developed saddle?

The saddle is usually a different colour to the rest of the body, and slightly wider


No → It is not a mature earthworm - you can't identify it with this Guide. At least 50% of the earthworms you find will be immature

Stripy earthworms

Which description best matches your worm? Is the body:

A Longer and wider or **B** Shorter and narrower?

A  Saddle usually pale

B  Saddle usually similar colour to the rest of the body

Hint
Can exude an unpleasant smelling yellowish fluid when handled

1. Compost worm *Eisenia veneta*

2. Brandling worm *Eisenia fetida*

Is the whole body clearly stripy on its upper surface when moving?


 **Yellow ring on body**

3. Green worm (green form) *Allolobophora chlorotica*

Hints
Often curls up in the hand
Yellow ring on body
Has 3 pairs of sucker-like discs (see 13)
Can exude a yellow fluid when handled

5. Black-headed worm


Aporrectodea longa

A 
Hint
Often a dark purplish head, the rear end of the body is much paler

B 
Hint
A stout worm, often as thick as a pencil

6. Lob worm *Lumbricus terrestris*

Sometimes it flattens its tail into a wide paddle shape


Are the male pores visible?


Is the earthworm longer than 8cm when NOT moving?
Red earthworms

Is the upper surface of the body, from the first segment to the saddle, entirely dark in colour (dark red, purplish red or chestnut brown)?


Is the body from the first segment to the saddle partly or entirely pale in colour (whitish, pink or grey)? It may have some reddish or dark segments


Hint
Sometimes slightly flattens its tail into a paddle shape


4. Redhead worm *Lumbricus rubellus*

Key continues on the other side

Return to start

Pale earthworms
Key continues on the other side

Photographs by: Harry Taylor¹ and Chris N. Lowe².
 Illustrations by: David T. Jones^{1,2}. Text by: David T. Jones^{1,2}, Chris N. Lowe², Harry Taylor², Paul Egglestone², Stephen Brooks², Emma Sherlock², Simon Norman⁴, Louise Parker⁴, Rebecca Farley¹, James Bone¹, Martin Head¹, Nick Vouliouklis¹, Linda Davies¹, Carolina Bachariou¹. ¹Imperial College London. ²Natural History Museum. ³Environment Agency. ⁴Field Studies Council. ⁵University of Central Lancashire. Supported by the Esmeé Fairbairn Foundation. Designed by FSC Publications.

© OPAL 2015. All rights reserved.



10. Rosy-tipped worm *Aporrectodea rosea*

Hints
 The head is usually rosy pink or pale pink up to the male pores
 Often has 2 or more whitish raised pads before the male pores. The saddle is usually orange, and can be wider towards the rear end

8. Chestnut worm *Lumbricus castaneus*

9. Little tree worm *Satchellius mammalis*

Is there a raised whitish gland visible on the underside between the saddle and the head?

Yes → [Chestnut worm]
No → [Next question]

Is the swelling around the male pore only covering one segment?

Yes → [Octagonal-tailed worm]
No → [Next question]

Are the male pores visible?

Visible → [Next question]
Not visible → [Red earthworms]

You have probably found a rare worm which is not in this key. Record it as unidentified

Is the swelling around the male pores broad, covering the entire width of adjacent segments?

Yes → [Little tree worm]
No → [Next question]

13. Green worm (pale form) *Allolobophora chlorotica*

Hints
 Has 3 pairs of sucker-like discs on the underside of alternate segments of the saddle, **not always easy to see**. The yellow ring can be faint
 Often curls up in the hand
 Can exude a yellow fluid when handled

Does the worm have this colour combination?

Pink or pale grey	Whitish	Darker grey
		Saddle

Yes → [Next question]
No → [Next question]

Does it have a yellow ring towards the head?

Yes → [Green worm]
No → [Next question]

Hints
 The front end up to the saddle is usually in three distinct shades: pink or pale grey, then whitish, and then darker grey
 The saddle pads usually form a two-humped ridge across three segments, but these can be difficult to see

11. Grey worm *Aporrectodea caliginosa*

12. Blue-grey worm *Octolasion cyaneum*

Hints
 Distinct yellow tail
 Can vary from faint blue-grey to a pale rosy pink colour
 May have a lilac-blue line on the upper surface

Tail not yellow or only slightly yellow

Are the last four or five segments distinctly yellow?

Yes → [Blue-grey worm]
No → [Next question]

