

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

CLINICAL STUDY PROTOCOL

Full Study Title:	Effect of Aspirin on Reducing Cancer & Improving Outcomes in Primary Sclerosing Cholangitis
Short Study title / Acronym:	Asp-PSC
Products:	Aspirin and Placebo
Development Phase:	Phase III
Sponsor:	Imperial College London
Version no:	3.0
Protocol Date:	18.02.2025
Funding	Cancer Research UK (CRUK, grant number PRCPJT-May22\100001)

Property of Imperial Clinical Trials Unit (ICTU)
May not be used, divulged or published without the consent of ICTU

This protocol has regard for the Health Research Authority (HRA) guidance

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

RESEARCH REFERENCE NUMBERS

IRAS ID:	1007320
REC Reference Number:	23/WA/0282
ISRCTN Number:	ISRCTN12358813
Sponsor ID:	23SM8165
Protocol Number:	C/45/2022
Funder reference:	PRCPJT-May22\100001

Keywords:

Primary Sclerosing Cholangitis, Aspirin, Randomised, Cancer

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

CONTACT LIST

Chief Investigators

Shahid A Khan
Hepatology and Gastroenterology Section
Division of Diabetes Endocrinology and Metabolism
Department of Medicine
Imperial College London
St Mary's Hospital Campus
South Wharf Road
W2 1NY
UK
Tel: +44 (0)203 312 6454/6254
Email: shahid.khan1@nhs.net

Simon Rushbrook
Norfolk and Norwich University Hospital
Colney Lane
Norwich
Norfolk
NR4 7UY
UK
Tel: 01603 641193
Email: simon.rushbrook@nnuh.nhs.uk

Co-Investigators

- Ruth Langley: Professor of Oncology & Clinical Trials, University College London
- Hassan Malik: Consultant Hepatobiliary Surgeon, Liverpool Hospital NHS Trust
- Nick Powell: Clinical Reader and Consultant in Gastroenterology, Imperial College London
- Palak Trivedi: Consultant Hepatologist and UK-PSC Steering Group Chief Investigator, University of Birmingham
- Catherine Williamson: Professor of Women's Health, Imperial College London (and honorary Professor of Women's Health, King's College London).

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

Sponsor

Research Governance and Integrity Team Imperial College London (RGIT)
Room 221, Level 2, Medical School Building
St Mary's Campus
Norfolk Place
London, W2 1PG
United Kingdom
Contact: Rinat Ezra
Clinical Trials Manager
Email: rgit.ctimp.team@imperial.ac.uk
Tel: +44 (0)20 7594 8081

Clinical queries

Clinical queries should be directed to the Chief Investigator or ICTU Study Manager who will direct the query to the appropriate person.

Funder

Cancer Research UK
2 Redman Place
London
E20 1JQ

ICTU Cancer

Meena Reddi
ICTU Study Manager
Imperial Clinical Trials Unit – Cancer
Cancer Research UK Convergence Science Centre
Department of Surgery and Cancer
Imperial College London
5th Floor Roderic Hill Building
South Kensington Campus
London
SW7 2AZ
Email: asp-psc@imperial.ac.uk

Joe Bolger
ICTU Trial Monitor
Imperial Clinical Trials Unit – Cancer
Cancer Research UK Convergence Science Centre
Department of Surgery and Cancer
Imperial College London

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

5th Floor Roderic Hill Building
South Kensington Campus
London
SW7 2AZ
Email: asp-psc@imperial.ac.uk

Lee Webber
ICTU Deputy Operations Manager
Imperial Clinical Trials Unit – Cancer
Cancer Research UK Convergence Science Centre
Department of Surgery and Cancer
Imperial College London
5th Floor Roderic Hill Building
South Kensington Campus
London
SW7 2AZ

Senior Statistician

Mr Nicholas Johnson
Research Fellow- Senior Statistician
Imperial Clinical Trials Unit (ICTU)
Imperial College London
Stadium House
68 Wood Lane
London W12 7RH
Email: nicholas.johnson@imperial.ac.uk

Central Storage of Laboratories

Norwich Research Park Biorepository
Norfolk and Norwich University Hospitals NHS Foundation Trust
Colney Lane
Norwich
NR4 7UY
Email: biorepository@nnuh.nhs.uk

iQur Laboratory

iQur Ltd
2 Royal College Street
NW1 0NH
Email: raakesh.modi@iqur.com

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

South West London Pathology

St George's University Hospitals NHS Foundation Trust
Blackshaw Rd, Tooting
London SW17 0QT

Black Country Pathology Services

Pathology Department
C37 Pathology Centre
New Cross Hospital
Wolverhampton
WV10 0QP

Cambridge University Hospitals

Pathology Directorate
Box 241
Cambridge University Hospitals
Hills Road
Cambridge
CB2 0QQ

Storage of Scans

Infinitt Europe GMBH
Somerset House
37 Temple Street
Birmingham
B2 5DP

IMP Manufacturing Facilities

Sabina Melander
Clinical Trial Manufacturing and Supplies
Royal Free Hospital Pharmacy
Royal Free Pharmacy Production Unit
Royal Free London NHS Foundation Trust
Royal Free Hospital
Pond Street
NW3 2QG
Tel:020 3758 2000 Ext. 22523
Email: sabina.melander@nhs.net

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

Protocol Development Group

Shahid A Khan Professor of Practice (Hepatology) and Chair Cholangiocarcinoma-UK	Imperial College London
Simon Rushbrook Consultant Hepatologist and UK-PSC Steering Group member	Norfolk and Norwich University Hospitals NHS Trust
Palak Trivedi Consultant Hepatologist and UK-PSC Steering Group Chief Investigator	University of Birmingham
Hassan Malik Consultant Hepatobiliary Surgeon and Chair of The British Association of Surgical Oncology (BASO)	Liverpool Hospital NHS Trust
Ruth Langley Professor of Oncology & Clinical Trials and Co-Chair, UK Therapeutic Cancer Prevention Network Group, (TCPN)	University College London
Nick Powell Clinical Reader and Consultant in Gastroenterology	Imperial College London
Catherine Williamson Professor of Women's Health	Imperial College London
Martine Walmsley Chair of PSC Support	PSC Support
Helen Morement Founder & Chief Executive Officer	Alan Morement Memorial Fund (AMMF)- The Cholangiocarcinoma Charity
Francesca Fiorentino Senior Lecturer in Clinical Trials Statistics	University of Leeds
Lee Webber ICTU Deputy Operations Manager	Imperial College London
Meena Reddi ICTU Study Manager	Imperial College London

This protocol describes the Asp-PSC trial and provides information about procedures for enrolling participants to the trial. The protocol should not be used as a guide for the treatment of other participants; every care was taken in its drafting, but corrections or amendments may be necessary. These will be circulated to investigators in the study, but centres enrolling participants for the first time are advised to contact ICTU to confirm they have the most recent version. Problems relating to this trial should be referred, in the first instance, to ICTU. This trial will adhere to the principles outlined in the Medicines for Human Use (Clinical Trials) Regulations 2004 (SI 2004/1031), amended regulations (SI 2006/1928) and the International Conference on Harmonisation Good Clinical Practice

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

(ICH GCP) guidelines. It will be conducted in compliance with the protocol, the Data Protection Act 2018 and other regulatory requirements as appropriate.

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

ABBREVIATIONS

AE	Adverse Event
AFP	Alpha-fetoprotein
AMMF	Alan Morement Memorial Fund
ASP	Aspirin
AST	Aspartate Aminotransferase
BASO	The British Association of Surgical Oncology
BMI	Body Mass Index
BTC	Biliary Tract Cancer
CA	Cancer Antigen
CCA	Cholangiocarcinoma
CI	Chief Investigator
CLDQ	Chronic Liver Disease Questionnaire
COX-2	Cyclooxygenase-2
CRC	Colorectal Cancer
CRF	Case Report Form
CRUK	Cancer Research United Kingdom
CRP	C-Reactive Protein
CT	Computerised Tomography
CTA	Clinical Trial Authorisation
CTIMP	Clinical Trial of an Investigational Medicinal Product
CTCAE	Common Terminology Criteria for Adverse Events
DMC	Data Monitoring Committee
DNA	Deoxyribonucleic acid
DSUR	Development Safety Update Report
eCCA	Extrahepatic Cholangiocarcinoma
eCRF	Electronic Case Report Form
ERCP	Endoscopic Retrograde Cholangiography
ELF	Enhanced Liver Fibrosis
FBC	Full Blood Count
GCP	Good Clinical Practice
GI	Gastrointestinal
GGT	Gamma-glutamyl Transferase

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

GMP	Good Manufacturing Practice
HR	Hazard Ratio
HCC	Hepatocellular Carcinoma
HPB	Hepato-Pancreato-Biliary
HRA	Health Research Authority
IBD	Inflammatory Bowel disease
ICMJE	International Committee of Medical Journal Editors
iCCA	Intrahepatic Cholangiocarcinoma
ICHNT	Imperial College Healthcare NHS Trust
ICMJE	International Committee of Medical Journal Editors
ICTU	Imperial Clinical Trials Unit
IDMC	Independent Data Monitoring Committee
IMP	Investigational Medicinal Product
IND	Investigational New Drug
INR	International Normalised Ratio
IRAS	Integrated Research Application System
ISRCTN	International Traditional Medicine Clinical Trial Registry
ITT	Intention to Treat
MDT	Multidisciplinary Team
MHRA	Medicines and Healthcare Products Regulatory Agency
MIA	Manufacturing Import Authorisation
MRCP	Magnetic Resonance Cholangiography
MRI	Magnetic Resonance Imaging
NASH	Non-alcoholic Steatohepatitis
NHS	National Health Service
NICE	National Institute for Health and Care Excellence
NSAID	Non-Steroidal Anti-Inflammatory drugs
OD	Once daily
PBC	Primary Biliary Cholangitis
PFIC	Progressive familial intrahepatic cholestasis
PPIE	Patient Public Involvement Engagement

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

PSC	Primary Sclerosing Cholangitis
PSC PRO	Primary Sclerosing Cholangitis Patient Reported Outcome
PTC	Percutaneous Transhepatic Cholangiogram
QA	Quality Assurance
QC	Quality Control
QOL	Quality of Life
REC	Research Ethics Committee
RGIT	Research Governance Integrity Team
RR	Risk ratio
RR	Relative risk
RSI	Reference Safety Information
SAE	Serious Adverse Event
SAR	Serious Adverse Reaction
SAP	Statistical Analysis Plan
SIBDQ	Short Inflammatory bowel disease questionnaire
SmPC	Summary of Product Characteristics
SF-36	Short Form-36
SOP	Standard Operating Procedure
SSAR	Suspected Serious Adverse Reaction
SUSAR	Suspected Unexpected Serious Adverse Reaction
TCPN	Therapeutic Cancer Prevention Network Group
TIA	Transient Ischaemic Attack
TMG	Trial Management Group
TSC	Trial Steering Committee
UDCA	Ursodeoxycholic Acid
UK	United Kingdom
US	Ultrasound
U & E	Urea and Electrolytes

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

1 Table of Contents

RESEARCH REFERENCE NUMBERS	2
CONTACT LIST	3
TRIAL SUMMARY	15
1 BACKGROUND	20
1.1 CLINICAL SETTING	20
1.1.1 Primary Sclerosing Cholangitis and Cancer Risk	20
1.2 INVESTIGATIONAL AGENT AND RATIONALE FOR STUDY	21
1.2.1 Evidence That Aspirin Imparts Protective Effects against CCA, CRC and HCC	21
1.3 POSTULATED ANTI-CANCER BIOLOGICAL MECHANISMS OF ASPIRIN	24
1.4 HYPOTHESIS	24
1.5 WHAT THE STUDY WILL ADD	24
1.6 RISK: BENEFIT RATIO	25
2 OBJECTIVES AND ENDPOINTS	26
2.1 PRIMARY OBJECTIVE	26
2.2 PRIMARY ENDPOINT	26
2.3 SECONDARY OBJECTIVE:	26
2.4 SECONDARY ENDPOINTS	26
2.5 TRANSLATIONAL RESEARCH OBJECTIVES:	26
2.6 TRANSLATIONAL RESEARCH OUTCOMES	27
2.7 SUB-STUDY OBJECTIVE:	27
2.8 ADDED BENEFIT:	27
3 STUDY DESIGN	27
3.1 DESIGN	27
3.2 TREATMENT REGIMENS	28
3.3 STUDY FLOW CHART SUMMARY	29
4 PARTICIPATION ENTRY	30
4.1 STUDY SETTING AND POPULATION	30
4.2 ANTICIPATED NUMBERS OF CENTRES AND EXPECTED ENROLMENT	31
4.3 INCLUSION/EXCLUSION CRITERIA	31
4.3.1 The Inclusion Criteria:	31
4.3.2 The Exclusion Criteria:	32
5 PROCEDURES AND MEASUREMENTS	33
5.1 IDENTIFICATION OF RECRUITMENT AND PARTICIPANTS	33
5.2 SCREENING AND PRE-RANDOMISATION EVALUATIONS	33
5.2.1 Screening assessments	33
5.3 RANDOMISATION AND BLINDING	34
5.4 CODE BREAKING/UNBLINDING	34
5.5 UNBLINDING	35
5.5.1 Gastrointestinal bleeding.....	35
5.5.2 Haemorrhagic Stroke / Intracranial Haemorrhage	35
5.5.3 Ischaemic Stroke / Transient Ischaemic Attack (TIA)	35
5.5.4 Surgery	36
5.6 VISIT SCHEDULE	36
5.7 FOLLOW-UP	38
5.7.1 Laboratory Evaluations	40
5.8 INCIDENTAL FINDINGS	41

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

6	TREATMENTS	42
6.1	TREATMENT ARMS	42
6.2	LABELLING AND PACKAGING	42
6.3	STORAGE AND DISPENSING	42
6.4	DOSAGE, DURATION AND COMPLIANCE	42
6.5	ACCOUNTABILITY	43
6.6	DRUG INTERACTIONS	43
6.6.1	Contraindicated Medications	43
6.6.2	Need to prescribe concurrent oral corticosteroids	43
6.7	DOSE MODIFICATIONS FOR ADVERSE EVENTS	43
6.8	OVERDOSE OF IMP	44
6.8.1	Symptoms	44
6.9	PERMANENT DISCONTINUATION OF STUDY TREATMENT	45
6.9.1	Withdrawal from Study	45
6.9.2	Procedures for Withdrawal from Study	46
6.9.3	Temporary Discontinuation of IMP in the study	46
7	PHARMACOVIGILANCE	46
7.1	ADVERSE EVENT (AE)	46
7.2	ADVERSE REACTION (AR)	46
7.3	UNEXPECTED ADVERSE REACTION	47
7.4	CAUSALITY	47
7.5	SEVERITY OF ADVERSE EVENTS	48
7.6	ADVERSE EVENT RECORDING	48
7.7	ABNORMAL LABORATORY TEST RESULTS	49
7.8	SERIOUS ADVERSE EVENTS (SAE)	49
7.8.1	Definition of SAE	49
7.9	REPORTING OF SAES	50
7.10	DEFINITION OF A SERIOUS ADVERSE REACTION (SAR)	50
7.10.1	Definition of Suspected Unexpected Serious Adverse Reaction (SUSAR)	50
7.10.2	Reporting of SUSARs	50
7.11	DEVELOPMENTAL SAFETY UPDATE REPORTS / ANNUAL SAFETY REPORTS	51
7.12	PREGNANCY	51
7.13	REPORTING URGENT SAFETY MEASURES	51
8	STATISTICAL ANALYSES	51
8.1	DESIGN	51
8.2	SAMPLE SIZE AND POWER CONSIDERATIONS	52
8.3	PLANNED RECRUITMENT RATE	52
8.4	STATISTICAL ANALYSIS	52
8.5	ANALYSIS OF PRIMARY AND SECONDARY OUTCOMES	52
8.6	PRIMARY EFFICACY OUTCOME ANALYSIS	52
8.7	SECONDARY OUTCOMES' ANALYSIS	53
8.8	MISSING DATA	53
9	REGULATORY, ETHICAL AND LEGAL ISSUE	53
9.1	DECLARATION OF HELSINKI	53
9.2	GOOD CLINICAL PRACTICE	53
9.3	RESEARCH ETHICS COMMITTEE (REC) APPROVAL	53

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

9.3.1	Initial Approval	53
9.3.2	Approval of Amendments	54
9.3.3	End of Trial Notification	54
9.4	REGULATORY AUTHORITY APPROVAL	54
9.5	HRA APPROVAL	54
9.6	NON-COMPLIANCE AND SERIOUS BREACHES	55
9.7	INSURANCE AND INDEMNITY AND SPONSOR	55
9.8	TRIAL REGISTRATION	55
9.9	INFORMED CONSENT	55
9.10	CONTACT WITH GENERAL PRACTITIONER	56
9.11	PARTICIPANT CONFIDENTIALITY	56
9.12	DATA PROTECTION AND PARTICIPANT CONFIDENTIALITY	56
9.13	END OF TRIAL	56
9.14	STUDY DOCUMENTATION AND DATA STORAGE	57
10	DATA MANAGEMENT	57
10.1	SOURCE DATA	57
10.2	LANGUAGE	57
10.3	DATABASE	57
10.4	DATA COLLECTION	58
10.5	ARCHIVING	58
11	STUDY MANAGEMENT STRUCTURE	58
11.1	TRIAL STEERING COMMITTEE	58
11.2	TRIAL MANAGEMENT GROUP	58
11.3	INDEPENDENT DATA MONITORING COMMITTEE	59
11.4	EARLY DISCONTINUATION OF THE STUDY	59
11.5	RISK ASSESSMENT	59
11.6	MONITORING	59
11.6.1	Central Monitoring at ICTU	59
11.6.2	On-site Monitoring	59
11.7	QUALITY CONTROL AND QUALITY ASSURANCE	60
11.8	PEER REVIEW	60
11.9	PUBLIC INVOLVEMENT	60
11.9.1	Patient and Public Involvement and Engagement (PPIE)	60
11.9.2	Patient & Public Involvement & Engagement (PPIE) Lead	61
11.9.3	PPIE Collaboration Plan	61
11.9.4	PPIE Evaluation and Feedback	62
11.10	PUBLICATION AND DISSEMINATION POLICY	62
12	REFERENCES	64
13	REVISION HISTORY	66
14	SIGNATURE PAGE 1 (CHIEF INVESTIGATORS)	68
15	SIGNATURE PAGE 2 (SPONSOR)	69
16	SIGNATURE PAGE 3 (STATISTICIAN)	70
17	SIGNATURE PAGE 4 (PRINCIPAL INVESTIGATOR)	71
18	APPENDICES	72

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

TRIAL SUMMARY

TITLE: Asp-PSC Trial: Effect of Aspirin on Reducing Cancer & Improving Outcomes in Primary Sclerosing Cholangitis (PSC)

OBJECTIVES:

Primary Objective:

To investigate if in patients with PSC and inflammatory bowel disease (IBD), daily low dose aspirin lowers the risk of PSC related cancer/high grade dysplasia, liver decompensation, the need for liver transplantation and all-cause mortality, compared to placebo, over a minimum 5-year period.

Secondary Objective:

Safety and tolerability of aspirin in patients with PSC-IBD and effect on quality of life using disease specific questionnaires.

Translational research objectives:

To collect, store and analyse clinical and radiological data, genomic deoxyribonucleic acid (DNA), serum, cell free DNA and urine samples.

Sub-study objective:

To assess safety, pregnancy outcome and acceptability of the intervention in pregnant women.

Added Benefit:

To create and develop pan-UK PSC bioresource for translational research to help improve the clinical care of patients with PSC.

PHASE: Asp-PSC is a Phase 3 study.

DESIGN: This study will be a multicentre, double-blind, randomised placebo-controlled trial. The study population will include patients with PSC who have a concurrent diagnosis of IBD and who are at least 12 months post diagnosis of PSC. Patients will be randomised in a ratio of 2:1 Aspirin vs Placebo. Accrual will be over 5 years, with a minimum participant follow up of 5 years. Included in this study is an embedded feasibility study, for one year from start of recruitment, to establish recruitment, acceptability, and compliance rates. Randomisation will be stratified by advanced fibrosis/cirrhosis, time since diagnosis of PSC and concomitant usage of oral 5-aminosalicylates.

SAMPLE SIZE: 774 patients need to be recruited to allow the detection of a hazard ratio (HR) of 0.6 with 80% power (significance 0.05), equivalent to a 6% difference in the proportion of patients diagnosed with cancer, undergoing liver transplantation or death from all-cause mortality. This is based on an estimated event rate in the control group of 15% at 5 years accrual and 5 years of minimum follow up. Of note, the number of individuals who develop hepatic decompensation specifically were not part of the original power calculation but are likely to be captured amongst those undergoing transplantation. The Intervention:

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

placebo randomisation ratio is 2:1, meaning we need to recruit 516 participants in the intervention group and 258 in placebo group. To allow for 20% loss to follow up, we have determined we will need to recruit 968 patients in total.

INCLUSION/EXCLUSION CRITERIA

Inclusion Criteria:

1. Age 18 years or above
2. Able to give written and informed consent.
3. Must have an established clinical diagnosis of large duct PSC-based on a standard disease definition of typical cholangiography findings on endoscopic retrograde cholangiography (ERCP) or magnetic resonance cholangiography (MRCP)
4. An established diagnosis of concomitant colonic IBD either in a pattern of Ulcerative Colitis, Crohn's disease or IBD unclassified.
5. Patients must be at least one-year post PSC diagnosis.
6. If pre-treated with ursodeoxycholic acid (UDCA) – UDCA therapy should remain at a stable dose for 12 weeks prior to screening, and not exceeding 20mg/kg/day.
7. Must have had a colonoscopy within the last year of the first day of screening date as part of routine clinical care. If not, this must be done within the screening interval.
8. If a patient has cirrhosis, they must have undertaken a hepatobiliary ultrasound (US), MRCP scan, magnetic resonance imaging (MRI) liver or regional computerised tomography (CT) scan within 6 months of screening date as part of routine clinical care. If not, this must be done within the screening interval.
9. If non-cirrhotic, then the patient must have had an US, MRCP, dynamic MRI or regional CT within the last 12 months as part of routine clinical care. If not, this must be done within the screening interval.

Exclusion Criteria

1. Evidence of concomitant disease(s) that causes secondary sclerosing cholangitis.
2. Evidence of any of the following diseases: IgG4 related disease, PBC, acute or chronic viral hepatitis, alcohol-related liver disease, Wilson disease, Budd-Chiari syndrome, portal vein thrombosis, alpha-1-antitrypsin disease, hepatic sarcoidosis, cystic fibrosis, Progressive familial intrahepatic cholestasis (PFIC), hereditary haemochromatosis, non-alcoholic steatohepatitis (NASH) – those with simple hepatic steatosis without evidence of NASH or liver fibrosis secondary to fatty liver disease are allowed to participate, other metabolic liver disease, active malignancy in the last five years (except treated/excised non-melanomatous skin cancer).
3. Have a previous diagnosis of colorectal cancer, cholangiocarcinoma, gallbladder cancer at any time point.
4. Has received a liver transplant, or is being actively assessed for liver transplant, or is listed for a liver transplant.
5. Had any of the following procedures: colonic resection of any nature, including those with a defunctioning colostomy.
6. Known diagnosis of alcohol related liver disease

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

7. Current or recent participation in any other clinical trial of an investigational medicinal product (CTIMP) within the last 6 weeks prior to first dose of aspirin/placebo
8. Already taking aspirin.
9. History of non-variceal upper gastrointestinal (GI) bleeding within one year.
10. A history of congestive cardiac failure.
11. A known diagnosis of glucose-6 -phosphate dehydrogenase.
12. Childs Pugh B or C cirrhosis
13. Known diagnosis of Lynch Syndrome or Li Fraumeni Syndrome
14. Vaccination for varicella zoster in the six weeks prior to screening period
15. Known allergy to aspirin
16. A history of NSAID/ aspirin induced asthma and nasal polyps
17. Chronic daily oral non-steroidal anti-inflammatory (NSAIDs) drug usage (continuously for 3 weeks prior to the first day of screening).
18. History of haemophilia and/or other bleeding disorders where aspirin is contraindicated
19. Taking other anti-platelet or anti-coagulant medication (e.g., clopidogrel, prasugrel, warfarin, apixaban, rivaroxaban, dabigatran or therapeutic heparin preparations)
20. Active, or history of recurrent peptic ulcer
21. Patients who are suffering from severe active gout
22. Severe renal impairment
23. Taking methotrexate at a dose of >15mg/week

TREATMENT/MAIN STUDY PROCEDURES

Patients will be randomised to either aspirin or placebo. Patients will be called at one month after the collection visit. The first month's call will be to ensure drug tolerability and check no immediate adverse events.

During the month one visit, the patients will be asked to for:

1. Medical History
2. Review of Haematology including FBR and INR test if locally available
3. Review of Biochemistry including UE, LFT, GGT, AST, CRP, Vitamin D, B12, ferritin, TSH and T4 and if locally available
4. Adverse Events
5. Concomitant Medications
6. QOL Assessments (PSC-PRO, SF-36, 5D-Itch Scale, SIBDQ, PRO-2 and CLDQ-PSC) to be completed at home.

During the one-month phone call patients may be required to attend an in person visit if clinically required to assess further adverse event reporting, this will be decided by a delegated member of staff responsible for patient care. The following assessments will be undertaken:

1. Vital Signs (Body temperature, pulse rate, respiration rate, blood pressure-if clinically indicated)
2. Targeted Physical Exam (if clinically indicated)

One-month questionnaires can be administered in the following ways:

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

- Given to the patient at the collection visit.
- Mailed to the patient's home in time for the one-month phone call.
- Administered over the phone by a designated staff member, who will record the patient's responses on a paper questionnaire. If this method is used, the staff member must note their name, signature, and date on the questionnaire.

Patients will be seen for study visits at 6 monthly intervals (5 months after the month 1 phone call, then 6 monthly thereafter), ideally in alignment with routine clinical care, for five years.

During each 6-month visit, the patients will be asked to:

1. Documentation of medical history
2. Documentation of vital signs (Body temperature, pulse rate, respiration rate, blood pressure)
3. Weight measurement
4. Targeted Physical Exam (if clinically indicated)
5. Undertake standard of care National Health Service (NHS) blood tests (Full Blood Count (FBC), Urea and Electrolytes (U&E), liver biochemistry, gamma-glutamyl transferase (GGT), aspartate aminotransferase (AST), C-Reactive Protein (CRP), International Normalised Ratio (INR), Vitamin D, Cancer Antigen 19.9 (CA19.9), Alpha-fetoprotein (AFP), ferritin and a TSH and T4). Standard of care faecal calprotectin measurement will be taken yearly.
6. Adverse event documentation
7. Complete questionnaires: Primary Sclerosing Cholangitis -Patient reported outcome (PSC PRO), Short Form-36 (SF-36), 5D-Itch Scale, and Short Inflammatory bowel disease questionnaire (SIBDQ), PRO-2 (Patient Reported Outcome-2) and Chronic Liver Disease Questionnaire (CLDQ)-PSC, questionnaires can be taken home to do. If Questionnaires are not brought to clinic, site will need to ask patient to complete questionnaires on the day of visit.
8. Documentation of Concomitant Medications
9. Urine and blood samples taken (if patient consented to this).
10. Body mass index (BMI) measurement recorded.
11. Patients will also be asked about their alcohol and smoking habits.
12. A fibroscan will also be undertaken if locally available.
13. Documentation of UKELD Score
14. Note will be made of any intervening cancer or high grade dysplasia diagnosis, decompensating liver events or referral for liver transplantation.
15. Note will also be made of any colonoscopy undertaken and also the histological findings from that examination. Note will also be made of any endoscopic intervention/ cholecystectomy or new cancer diagnosis made.
16. Note will also be made of any recent imaging studies the patients have undergone, the results available from these studies and multidisciplinary team (MDT) outcomes.
17. Collection of aspirin/placebo tablets
18. Return of the one month questionnaire (if completed at home)

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

Patients will attend an end of treatment visit, 28 days after last study dose.

Patients will then be followed up yearly for a minimum of 5 years for data collection only.

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

OUTCOME MEASURES

PRIMARY ENDPOINT

The composite singular primary endpoint is defined as the occurrence of any of the following:

1. Hepatobiliary cancer (including gallbladder cancer/high grade dysplasia, pancreas, cholangiocarcinoma [CCA] or hepatocellular [HCC]) or colorectal cancer/high grade dysplasia
2. Listing for Liver transplantation
3. All-cause mortality

SECONDARY ENDPOINTS

1. To investigate gastrointestinal bleeding risk ((Common Terminology Criteria for Adverse Events (CTCAE) grade 3 or 4)) between daily low dose aspirin and placebo in patients with PSC-IBD
2. Progressive liver disease, as evidenced by either ascites, variceal bleeding (any site), hepatic encephalopathy (investigator discretion) or development of cirrhosis with Child Pugh B or C stage
3. To investigate the rates of referral for liver transplantation between daily low dose aspirin and placebo in patients with PSC-IBD
4. To investigate the rates of acute cholangitis between daily low dose aspirin and placebo in patients with PSC-IBD
5. To investigate IBD flare frequency between daily low dose aspirin and placebo in patients with PSC-IBD
6. To investigate the rates colonic surgery/resection for severe IBD between daily low dose aspirin and placebo in patients with PSC-IBD
7. Safety and tolerability of aspirin in patients with PSC-IBD and effect on quality of life using disease specific questionnaires.

TRANSLATIONAL RESEARCH OUTCOMES

To develop a prospective clinical, imaging and specimen biobank of samples from patients with PSC-IBD to allow future studies to be undertaken in which the performance of novel diagnostic biomarkers for the early detection and prognostication of PSC related cancer will be determined.

IMP: 75mg Aspirin or matched placebo taken orally daily once daily (OD) for five years.

1 BACKGROUND

1.1 CLINICAL SETTING

1.1.1 Primary Sclerosing Cholangitis and Cancer Risk

PSC is an immune-mediated liver disease which is associated with a unique type of IBD of the colon. Chronic inflammation at biliary and colonic mucosal sites results in the potential for epithelial cells to undergo a dysplasia/adenocarcinoma transition sequence (1, 2). No

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

medical therapy has been proven to slow disease progression. Consequently, over 50% of patients develop hepatopancreatobiliary (HPB) cancer or chronic liver failure in need of transplantation. Moreover, for the majority of patients (around 70%) that develop IBD there is a 30% (21) lifetime risk of colorectal cancer (CRC) – tenfold greater than that of IBD alone. The estimated prevalence of PSC in the UK is around 6/100,000 and PSC represents over 10% of all UK indications for liver organ transplantation. Moreover, PSC is now leading indication for liver transplantation in several European countries (3, 4).

The standardised incidence ratio of HPB cancer and CRC in PSC is estimated at around 400-fold (5); and a recent systematic review and meta-analysis reported summary relative risks (RR) of 584.4 (CI 269.4 – 1267.5) for CCA, 155.5 (125.3 – 193.0) for HPB cancer overall, 30.2 (12.0 – 76.2) for liver cancer including HCC, 6.1 (4.2 – 8.9) for colorectal cancer (CRC) and 4.1 (3.0 – 5.7) for all cancers (6). Furthermore, for those that develop a bile duct cancer, the overall survival is normally less than one-year from diagnosis.

PSC is also the commonest predisposing cause of CCA, itself the second commonest primary liver cancer in the Western World (7). The risk for developing CCA in PSC is highest within the first year of diagnosis (8). After that, the yearly incidence of CCA is 0.5-1.5%, with a lifetime risk of 15-18%, and a risk ratio (RR) of 150 times that of the general population.

At present due to a lack of evidence, there is no agreed guidance on how surveillance for CCA should be undertaken for patients with PSC, nor consistency in published national or international guidelines. In addition, a recent large-scale audit of clinical care showed a complete lack of a co-ordinated approach in the UK for cancer surveillance in PSC with some not offering it at all (9). This is despite a wealth of evidence indicating improved post-cancer survivorship for patients undergoing annual cancer surveillance with combination HPB imaging (10). However, the optimal imaging modality and interval for surveillance is unknown.

Patients with PSC and HPB cancer frequently present at a late and incurable stage of disease with an average life expectancy being less than 12 months from diagnosis. Moreover, whilst patients with PSC-induced decompensated liver disease may be eligible for liver transplantation, the development of cancer is currently a contraindication in the UK. In the few countries where liver transplantation for CCA is carried out, only a very few per cent of PSC patients qualify (7). ***There is, therefore, an urgent and unmet need to find new therapies that could prolong cancer free survival in PSC.***

1.2 INVESTIGATIONAL AGENT AND RATIONALE FOR STUDY

1.2.1 Evidence That Aspirin Imparts Protective Effects against CCA, CRC and HCC

Based on pre-clinical, epidemiological and randomised data aspirin may have anti-cancer effects, with the strongest evidence currently pertaining to CRC.

Cohort studies found that aspirin significantly reduces the risk of CRC in the general population. In a 2017 nested case-control study set in UK primary care (11), users of low-dose aspirin had a significantly reduced risk of CRC (RR 0.66, 95% CI 0.60-0.74). The reduction in risk was apparent across all age strata and with all aspirin groups. After five years of therapy, the RR for Dukes Stage A CRC was 0.53 (0.24-1.19). In a separate

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

retrospective cohort study of over 612,000 patients, 5,118 (2.5%) out of 204,170 aspirin users were diagnosed with CRC; and 2073 (1.0%) died of malignancy. Aspirin usage significantly reduced CRC mortality (HR = 0.59; 95% CI 0.56 to 0.62) (12).

In the double-blind, randomised CAPP2 trial, 861 patients with Lynch syndrome, which carries a high risk of CRC, patients were randomly assigned to receive aspirin daily or placebo. The long-term results of the CAPP2 trial found, on a per-protocol analysis, significantly reduced overall risk for the aspirin group: HR 0.63 (CI 0.43-0.92; $p=0.018$) (13). The recommendation that these patients be considered for daily aspirin has recently been incorporated into National Institute for Health and Care Excellence (NICE) guidelines (14).

Epidemiological evidence has also been accumulating in recent years that aspirin use is associated with a reduced risk of developing hepatobiliary cancer, including in PSC. A recent meta-analysis of all observational studies on aspirin for the prevention of digestive tract cancers estimated the pooled relative risk (RR) of cancer for regular aspirin use versus non-use using random effects models (15). In this study, regular aspirin use was associated with a reduced risk of crc (RR 0.73, 95% CI 0.69-0.78, 45 studies), squamous-cell oesophageal cancer (RR 0.67, 95% CI 0.57-0.79, 13 studies), adenocarcinoma of the oesophagus and gastric cardia (RR 0.61, 95% CI 0.49 – 0.77, 10 studies), stomach cancer (RR 0.64, 95% CI 0.51-0.82, 14 studies), hepato-biliary tract cancer (RR 0.62, 95% CI 0.440-0.86, five studies), and pancreatic cancer (RR 0.78, 95% CI 0.68-0.89, 15 studies).

Focussing on CCA, a hospital-based case-control study found that aspirin use was significantly associated with a 2.7 to 3.6-fold decreased risk for both main anatomical subtypes of CCA: intrahepatic (iCCA) and extrahepatic (eCCA), with adjusted odds ratios (OR) of 0.29 – 0.35 (16). Furthermore, there was up to a 50% reduction specifically in PSC-associated CCA. Further support for aspirin's potentially protective effect against CCA in the wider population comes from two meta-analyses. Lapummuaypol et al.'s systematic review examined five observational studies with a total of over 9.2 million enrolled patients (17). The pooled OR of CCA in aspirin users was 0.56. No significant association between NSAID use and CCA was found.

Additionally, a systematic review and meta-analysis by Xiong et al. examined nine studies, consisting of 12,535 people who developed CCA and 92,750 healthy controls (18). There was a significantly decreased risk of CCA in those using aspirin. Moreover, this relationship was detected only in case-control studies (OR=0.65; 95% CI=0.38–0.93), and the reduction was across all anatomical sites of CCA (iCCA, OR=0.33 (95% CI=0.26–0.39) & eCCA, OR=0.56, 95% CI=0.41–0.73).

There is also emerging epidemiological evidence that aspirin may have benefit as adjuvant therapy (19), including improving survival after a diagnosis of biliary tract cancer (BTC). Jackson et al. investigated the association between post-diagnosis aspirin use and BTC survival (20). Even use of post-diagnosis aspirin, at low dose, was associated with decreased risk of death in patients with Gall Bladder Cancer (HR, 0.63; 95% CI, 0.48-0.83), CCA (HR, 0.71; 95% CI, 0.60-0.85), Ampulla of Vater Cancer (HR, 0.44; 95% CI, 0.26-0.76), and overlapping BTC (HR, 0.68; 95% CI, 0.50-0.92). The authors noted that the survival benefit of aspirin observed in their study is on par with the current standard of care for these cancers.

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

This work is further supported by Liao et al.'s recent study of newly diagnosed BTC patients included 2,519 (16%) taking aspirin post BTC diagnosis (21). After a mean follow-up of 1.59 years, the 5-year survival rate was 27.4%. The multivariate-adjusted HR for postdiagnosis aspirin users, as compared with nonusers, was 0.55 (95% CI: 0.51 to 0.58) for BTC-specific death. Cancer-specific mortality was lower with post-diagnosis aspirin use in patients with all major BTC subtypes. The authors concluded randomised trials are required to investigate aspirin's efficacy in BTC.

A recent nationwide Swedish registry study examined long term effects of low dose aspirin on incident HCC, liver related mortality and gastrointestinal bleeding in persons with chronic viral hepatitis B or C infection, which are established risk factors for cirrhosis, HCC and CCA (22). Over 50,000 non-aspirin users were compared to over 14,000 patients taking low dose aspirin. Use of aspirin was associated with a significantly lower risk of HCC than no use of aspirin. Adjusted HRs were 0.90 (95% CI, 0.76 to 1.06) for 1 to less than 3 years of aspirin use, 0.66 (95% CI, 0.56 to 0.78) for 3 to less than 5 years of use, and 0.57 (95% CI, 0.42 to 0.70) for 5 or more years of use. The 10-year risk of gastrointestinal bleeding did not differ significantly between users and nonusers of aspirin.

A summary of the statistically significant findings of the protective effects of aspirin use against HPB cancers and CRC is provided in Table 1.

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

Table 1: Summary of studies reporting statistically significant protective effects of Aspirin against cancers PSC-IBD patients are at high risk of

Evidence of Aspirin Users' Potential Protection against HPB and CRC Cancers
1. Choi et al. 2016: OR 0.29 – 0.35 for all CCA subtypes (50% reduction in PSC associated CCA) (16)
2. Simon et al. 2020: HR 0.57 for HCC (22)
3. Lapumnuaypol et al. 2019: OR 0.56 for CCA (Meta-analysis) (17)
4. Xiong et al. 2018: iCCA OR 0.33 for iCCA; OR=0.56 for eCCA (Meta-analysis) (18)
5. Bosetti et al. 2020: RR 0.62 for HPB cancers (15)
6. Garcia Rodriguez et al. 2017: RR 0.53 for Dukes Stage A CRC (11)
7. Burn et al 2020: CAPP-2 Randomised Controlled Trial: HR 0.63 for CRC in Lynch syndrome (13)
Post-diagnosis aspirin use associated with decreased biliary tract cancer-specific mortality
7. Liao et al. 2021: HR 0.55 (21)
8. Jackson et al. 2019: HR 0.44 to 0.71 across all BTC sites (20)

1.3 POSTULATED ANTI-CANCER BIOLOGICAL MECHANISMS OF ASPIRIN

The biological plausibility of aspirin reducing the risk of CCA and CRC, in parallel to the published epidemiological data, has been extensively reviewed and includes its ability to: inhibit Cyclooxygenase-2 (COX-2) in dysplastic tissue (which promotes inflammation and cell proliferation), inhibit activation of nuclear factor Kb, prevent platelet aggregation (which may slow metastatic spread), modulate Wnt signalling via both COX-dependent and COX-independent pathways, upregulate tumour suppressor genes and stabilise DNA mismatch-repair proteins (23). Furthermore, established and recognised cell lines of CCA have been shown to be inhibited by aspirin (24). Therefore, given the unmet need for patients with PSC and also the clear epidemiological evidence that aspirin reduces cancers that reduce survival in PSC we have proposed the following hypothesis that this protocol will test.

1.4 HYPOTHESIS

Aspirin reduces the risk of PSC-related cancer occurrence, prolongs overall survival and liver transplant free survival in patients with both PSC-IBD.

1.5 WHAT THE STUDY WILL ADD

We anticipate this study will show that low dose daily aspirin safely results in improvements in cancer free survival, overall survival and liver transplant free survival in

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

patients with PSC-IBD. If so, this would lead to a change in clinical practice nationally and internationally; and will positively address a major area of unmet need in this cohort of patients, for whom no therapy currently has been shown to alter the natural history of disease progression and cancer free survival. A positive result in favour of aspirin would trigger a rapid review and update of the British Society of Gastroenterology’s UK-PSC Clinical Practice Guidelines (1) and we anticipate that a new recommendation to repurpose daily low-dose aspirin (a readily available, low-cost medication) to people with PSC would mean our findings could be put into practice almost immediately after the end of the trial.

1.6 RISK: BENEFIT RATIO

The overall risks of low dose aspirin are deemed to be low, based on several published trials of aspirin use in cancer prevention, often at much higher doses than we will be using. The safety of aspirin is key to several of our secondary endpoints. The potential benefits from the literature include lower cancer risk and improved survival.

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

2 OBJECTIVES AND ENDPOINTS

2.1 PRIMARY OBJECTIVE

To investigate if in patients with PSC and inflammatory bowel disease (IBD), daily low dose aspirin lowers the risk of PSC related cancer/high grade dysplasia, liver decompensation, the need for liver transplantation and all-cause mortality, compared to placebo, over a minimum 5-year period.

2.2 PRIMARY ENDPOINT

The composite singular primary endpoint is defined as occurrence of any of the following:

- Hepatobiliary cancer (including gallbladder cancer/high grade dysplasia, pancreas, cholangiocarcinoma [CCA] or hepatocellular [HCC]) or colorectal cancer/high grade dysplasia
- Listing for Liver transplantation
- All-cause mortality

2.3 SECONDARY OBJECTIVE:

Safety and tolerability of aspirin in patients with PSC-IBD and effect on quality of life using disease specific questionnaires.

2.4 SECONDARY ENDPOINTS

1. To investigate gastrointestinal bleeding risk ((Common Terminology Criteria for Adverse Events (CTCAE) grade 3 or 4)) between daily low dose aspirin and placebo in patients with PSC-IBD
2. Progressive liver disease, as evidenced by either ascites, variceal bleeding (any site), hepatic encephalopathy (investigator discretion) or development of cirrhosis with Child Pugh B or C stage
3. To investigate the rates of referral for liver transplantation between daily low dose aspirin and placebo in patients with PSC-IBD
4. To investigate the rates of acute cholangitis between daily low dose aspirin and placebo in patients with PSC-IBD
5. To investigate IBD flare frequency between daily low dose aspirin and placebo in patients with PSC-IBD
6. To investigate the rates colonic surgery/resection for severe IBD between daily low dose aspirin and placebo in patients with PSC-IBD
7. To investigate the quality of life between daily low dose aspirin and placebo in patients with PSC-IBD using disease specific questionnaires.

2.5 TRANSLATIONAL RESEARCH OBJECTIVES:

To collect, store and analyse clinical and radiological data, genomic deoxyribonucleic acid (DNA), serum, cell free DNA and urine samples.

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

2.6 TRANSLATIONAL RESEARCH OUTCOMES

To develop a prospective clinical, imaging and specimen biobank of samples from patients with PSC-IBD to allow future studies to be undertaken in which the performance of novel diagnostic biomarkers for the early detection and prognostication of PSC related cancer will be determined.

2.7 SUB-STUDY OBJECTIVE:

To assess safety, pregnancy outcome and acceptability of the intervention in pregnant women.

2.8 ADDED BENEFIT:

To create and develop pan-UK PSC bioresource for translational research to help improve the clinical care of patients with PSC.

The timepoint of evaluation of all endpoints are over a minimum period of 5 years of follow up. Please see table in section 5.6.

3 STUDY DESIGN

This study will be performed at approximately 60 investigational sites in the United Kingdom. This is a randomised, double-blind, multicentre, placebo-controlled study performed with a 2:1 randomisation ratio of Aspirin vs placebo. Participants will be randomised to one of 2 treatments, as shown in Table 3.2. Aspirin or placebo will be administered in single oral dose of 75 mg once per day for 5 years.

3.1 DESIGN

The Asp-PSC study will be a prospective, multicentre, Phase III double-blind, randomised controlled trial across the UK. **The intervention will be 75mg OD of aspirin** versus placebo. The study will be coordinated by the ICTU. Patients that have a diagnosis of large duct PSC of at least greater than one year in duration and IBD will be invited to take part. For the purpose of this study the time of diagnosis will be from when they had a confirmatory diagnostic investigation i.e., MRCP, ERCP, percutaneous transhepatic cholangiogram (PTC) +/- liver biopsy.

The study will randomise participants in a 2:1 ratio of aspirin vs placebo. The choice of randomisation ratio design was guided by our Patient and Public Involvement and Engagement (PPIE) group. The PPIE group advocated that such design would increase interest in participation in the study, as the probability of getting the active intervention (low dose aspirin) is higher than getting the placebo.

Recruitment will be completed within 5 years, with a minimum required participant follow up of 5 years. An embedded feasibility phase will be included to establish recruitment and acceptability and determine continuation of funding.

Randomisation will be stratified by fibrosis stage (as determined by Enhanced Liver Fibrosis (ELF) markers and/or fibroscan (if ELF test cannot be performed); time (in years)

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------


since PSC diagnosis (as determined by 1 - <5 , ≥5 - <10, ≥10), and concomitant aminosalicylate use.

3.2 TREATMENT REGIMENS

Treatments	Estimated number of <i>participants</i> needed to detect the predefined effect size	Number of participants to be recruited including a 20% drop out
Aspirin	516	645
Placebo	258	323
Total number of <i>participants</i>	774	968

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

3.3 STUDY FLOW CHART SUMMARY



Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

4 PARTICIPATION ENTRY

4.1 STUDY SETTING AND POPULATION

The patients proposed in this study will be enrolled from NHS hospitals trusts in the UK who form part of the established NIHR Liver Clinical Research Network BASL-SIG for Immune-Mediated Liver Disease. The Asp-PSC study will be done in conjunction/parallel with the prospective observational UK-PSC registry study to maximise patient recruitment.

UK-PSC was initially a large NIHR study set up by Dr Rushbrook in 2008 to look at the genetics of PSC. In this study, UK-PSC established a pan-UK patient cohort that led to several large genetic studies being undertaken in PSC along with high level clinical phenotyping data. Over 2,900 PSC patients have so far consented and been recruited to this project. This covers over 190 NHS trusts. We intend to directly approach PSC leads where these patients are seen. We anticipate at least 30% of these Trusts will recruit patients to this trial and therefore plan to have 60 active sites. To support recruitment, PSC Support (the UK patient organisation for people affected by PSC) will share information about the study to the PSC community through their dedicated research news bulletins and online patient support group of 3,300 members.

In the set-up phase of this trial the investigator team will continue the already begun series of preliminary meetings with recognised UK PSC investigators from various NHS sites from around the UK. Following on from this, the Asp-PSC study managerial team will set up the study with the local Research and Development team. Given the nature of this trial we will aim to do this across 60 NHS trusts, around 40% of the sites that have recruited to the UK-PSC network. Table 4.2 summarises the anticipated numbers of centres, and expected enrolment, and Figure 2 graphs the projected recruitment over 5 years. Patients recruited in the first year's feasibility stage (see section 4.2) will be included in the final analysis.

The NIHR BioResource is a recallable resource of over 300,000 volunteers, with and without health conditions who have agreed to be contacted about taking part in health-related research. We will be recruiting from within the IBD BioResource which forms part of the larger NIHR BioResource. Participants in the IBD BioResource have been recruited through clinics and have a confirmed diagnosis of IBD. NIHR BioResource, under their ethical approval, will identify participants from their database who are potentially eligible for the study. Staff at NIHR BioResource have ethical approval to send out invites to these participants, which summarise the study, what's involved, next steps, how they can take part, and they also include Participant Information Sheets. Once a participant expresses an interest in taking part their information will be passed onto the research team. They will then contact the participant and if relevant their treating clinician and arrange for screening. The research team will inform the BioResource whether or not the participant takes part in and this will be recorded on the BioResource database as per their protocols.

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

4.2 ANTICIPATED NUMBERS OF CENTRES AND EXPECTED ENROLMENT

Year	Number of Centres (cumulative)	Type and number of Centres opened	No. patients enrolled (cumulative)
1	10	Transplant Centres (N=5) Tier 2 Liver Centres (N=5)	80
2	30	Transplant Centres (N=5) Tier 2 Liver Centres (N=10) Other Centres (N=15)	300
3	40	Transplant Centres (N=5) Tier 2 Liver Centres (N=15) Other Centres (N=20)	500
4	60	Transplant Centres (N=5) Tier 2 Liver Centres (N=20) Other Centres (N=35)	750
5	60	Transplant Centres (N=5) Tier 2 Liver Centres (N=20) Other Centres (N=35)	968

There will be **no exceptions** (waivers) to eligibility requirements at the time of enrolment. Questions about eligibility criteria should be addressed **prior** to attempting to enrol the participant.

The eligibility criteria for this trial have been carefully considered and are the standards used to ensure that only medically appropriate participants are entered. Participants not meeting the criteria should not be entered into the trial for their safety and to ensure that the trial results can be appropriately used to make future treatment decisions for other people with similar diseases or conditions. It is therefore vital that exceptions are not made to these eligibility criteria.

Participants will be considered eligible for enrolment in this trial if they fulfil all the inclusion criteria and none of the exclusion criteria as defined below.

4.3 INCLUSION/EXCLUSION CRITERIA

4.3.1 The Inclusion Criteria:

1. Age 18 years or above
2. Able to give written and informed consent.
3. Must have an established clinical diagnosis of large duct PSC-based on a standard disease definition of typical cholangiography findings on endoscopic retrograde cholangiography (ERCP) or magnetic resonance cholangiography (MRCP).
4. An established diagnosis of concomitant colonic IBD either in a pattern of Ulcerative Colitis, Crohn's disease or IBD unclassified.
5. Patients must be at least one-year post PSC diagnosis.
6. If pre-treated with ursodeoxycholic acid (UDCA) – UDCA therapy should remain at a stable dose for 12 weeks prior to screening, and not exceeding 20mg/kg/day.

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

7. Must have had a colonoscopy within the last year of the first day of screening date as part of routine clinical care. If not, this must be done within the screening interval.
8. If a patient has cirrhosis, they must have undertaken a hepatobiliary ultrasound (US), MRCP scan, magnetic resonance imaging (MRI) liver or regional computerised tomography (CT) scan within 6 months of screening date as part of routine clinical care. If not, this must be done within the screening interval.
9. If non-cirrhotic, then the patient must have had an US, MRCP, dynamic MRI or regional CT within the last 12 months as part of routine clinical care. If not, this must be done within the screening interval.

4.3.2 The Exclusion Criteria:

1. Evidence of concomitant disease(s) that causes secondary sclerosing cholangitis.
2. Evidence of any of the following diseases: IgG4 related disease, PBC, acute or chronic viral hepatitis, alcohol-related liver disease, Wilson disease, Budd-Chiari syndrome, portal vein thrombosis, alpha-1-antitrypsin disease, hepatic sarcoidosis, cystic fibrosis, Progressive familial intrahepatic cholestasis (PFIC), hereditary haemochromatosis, non-alcoholic steatohepatitis (NASH) – those with simple hepatic steatosis without evidence of NASH or liver fibrosis secondary to fatty liver disease are allowed to participate, other metabolic liver disease, active malignancy in the last five years (except treated/excised non-melanomatous skin cancer).
3. Have a previous diagnosis of colorectal cancer, cholangiocarcinoma, gallbladder cancer at any time point.
4. Has received a liver transplant, or has been actively assessed for liver transplant, or is listed for a liver transplant.
5. Had any of the following procedures: colonic resection of any nature, including those with a defunctioning colostomy.
6. Known diagnosis of alcohol related liver disease
7. Current or recent participation in any other clinical trial of an investigational medicinal product (CTIMP) within the last 6 weeks prior to first dose of aspirin/placebo
8. Already taking aspirin.
9. History of non-variceal upper gastrointestinal (GI) bleeding within one year.
10. A history of congestive cardiac failure.
11. A known diagnosis of glucose-6 -phosphate dehydrogenase.
12. Childs Pugh B or C cirrhosis
13. Known diagnosis of Lynch Syndrome or Li Fraumeni Syndrome .
14. Vaccination for varicella zoster in the six weeks prior to screening period
15. Known allergy to aspirin
16. A history of NSAID/ aspirin induced asthma and nasal polyps
17. Chronic daily oral non-steroidal anti-inflammatory drugs (NSAIDs) drug usage (continuously for 3 weeks prior to the first day of screening)
18. History of haemophilia and/or other bleeding disorders where aspirin is contraindicated
19. Taking other anti-platelet or anti-coagulant medication (e.g., clopidogrel, prasugrel, warfarin, apixaban, rivaroxaban, dabigatran or therapeutic heparin preparations)
20. Active, or history of recurrent peptic ulcer
21. Patients who are suffering from severe active gout
22. Severe renal impairment
23. Taking methotrexate at a dose of >15mg/week

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

5 PROCEDURES AND MEASUREMENTS

5.1 IDENTIFICATION OF RECRUITMENT AND PARTICIPANTS

Patients will be identified by the local PI who will be responsible for the study conduct at each site and by their associated investigator team as determined by the local study delegation log.

Patients will be identified by means of the following ways:

- 1) From local clinic lists
- 2) From lists of patients identified by the British Association of the Study of the liver/BSG PSC UK Audit
- 3) From lists of patients held by UK-PSC for the local site.
- 4) From clinics in a prospective manner
- 5) The NIHR BioResource (see section 4)

Local PIs and the central team will have ethical permission to write to patients to invite them to take part in the study locally. Once randomised patients (following signed consent and undertaking a defined 60-day screening period) will be called at one month following collection visit. This study visit will be to ensure drug tolerability and check no immediate adverse events. After this time point, patients will be seen at 6 monthly intervals (5 months after month 1 visit, then 6 monthly thereafter) in alignment with routine clinical care during which a research visit will be undertaken.

5.2 SCREENING AND PRE-RANDOMISATION EVALUATIONS

Written informed consent to enter and be randomised into the trial must be obtained from participants after explanation of the aims, methods, benefits and potential hazards of the trial and **before** any trial-specific procedures are undertaken. The only procedures that may be performed in advance of written informed consent being obtained are those that would be performed on all patients as a usual standard of care e.g., annual colonoscopy for IBD, fibroscan, liver imaging and routine biochemical or haematological analysis.

5.2.1 Screening assessments

1. Confirmation the patient has consented.
2. Confirmation that the patient meets both the inclusion and exclusion criteria.
3. Confirmation that the patient has had a colonoscopy within the last 12 months of the screening date with either random biopsies or dye surveillance.
4. Confirmation that if the patient is cirrhotic (defined by either liver histology, imaging features, or vibration controlled transient elastography (VCTE) (KPa >14.4)), the patient has a liver USS or CT/dynamic MRI/MRCP within the last 6 months.
5. Confirmation that if the patient is non-cirrhotic, they have undergone a form of liver imaging (either US, MRI liver with contrast, MRCP or regional CT) with in the last year of randomisation date.

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

6. Determine of liver fibrosis by Fibroscan (if ELF test cannot be performed) or ELF testing.
7. Documentation of past medical history
8. Documentation of prescribed medication
9. Documentation of demographics
10. Documentation of vital signs
11. Documentation of height and weight
12. Documentation of adverse events
13. Questionnaires: PSC-PRO, SF-36, 5D-Itch Scale, and SIBDQ, PRO-2 and PSC CLDQ-PSC
14. Routine NHS blood tests (FBC, UE, LFT, GGT, AST, CRP, INR, Vitamin D, CA19.9, AFP, ferritin and a TSH and T4
15. Physical examination
16. Research blood and urine samples if patient consented.
17. ERCP/ Spyglass Cholangioscopy result collection if locally available.

5.3 RANDOMISATION AND BLINDING

In this study patients will be randomised to aspirin or placebo in a 2:1 ratio, in a double blinded manner. Within this study a small number of factors have been preidentified that will need to be controlled for in the randomisation process. We therefore utilise randomisation stratified by degree of fibrosis (binary, as determined by ELF score >9.8), OR Fibroscan score (If ELF test cannot be performed) of >9.6 Kpa), PSC disease duration (in years, as determined by 1 - <5 , ≥5 - <10, ≥10) and concurrent use of aminosalicylates (yes or no). Once the patient has been randomised, they will be asked to attend a collection visit for collection of aspirin/placebo.

5.4 CODE BREAKING/UNBLINDING

The investigator has the primary right to break blinding to treat participant in emergency situations and have access for rapid unblinding. The opinion to break the blind is with the investigator, and that the investigator does not need to discuss with the sponsor.

If a local investigator is required to unblind the patient, this can be done within the sealed envelope system in case of an emergency with additional help provided by the ICTU Study Manager or ICTU Trials Monitor. Unblinding should generally only be considered in the event of medical or safety needs (for example, an operation is required, or an overdose has been taken) where knowledge of the participant's treatment allocation could change clinical management. Following unblinding, the reason for the decision to unblind and the relevant parties must be documented in the participant's medical record. Treatment identification information should be kept confidential and should be disseminated only to those individuals that must be informed for medical management of the participant. If a participant requires unblinding, for example due to an upper GI bleed, they will have reached an end point and come off IMP but enter yearly follow up, see withdrawal section (6.9.1).. Their data will be used in the final analysis. An unblinding manual will be provided to sites to instruct on the unblinding process.

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

5.5 UNBLINDING

In most acute clinical situations, it should be possible to clinically manage a participant by assuming they have been taking 75mg of aspirin daily and/or by subsequently withholding the trial medication if appropriate. Nonetheless, there may be occasional clinical circumstances where knowledge of treatment allocation is essential to help guide urgent clinical management. If unblinding is considered, this should first be discussed with the Trials Unit and/or the local trial Principal Investigator wherever possible.

5.5.1 Gastrointestinal bleeding

Participants suspected of having a gastrointestinal bleed should have their trial medication stopped and should be managed according to standard local pathways. Standard treatment for gastrointestinal bleed should be initiated without delay and the patient should be managed by assuming they have been taking aspirin 75mg daily. It is not expected that emergency unblinding should be necessary in such events. Nonetheless, if a situation arises where knowledge of the trial treatment allocation would alter the immediate clinical management of a participant, this should be confirmed by the relevant specialist locally and unblinding can be performed.

5.5.2 Haemorrhagic Stroke / Intracranial Haemorrhage

Participants suspected of having an intracranial haemorrhage should have their trial medication withheld and should undergo appropriate neuroimaging and clinical management without delay. If the diagnosis is confirmed, then the trial medication should be stopped. It is not expected that emergency unblinding will be necessary in the event of an intracranial haemorrhage, however if a situation arises where knowledge of the trial treatment allocation would alter the immediate clinical management of a participant, this should be confirmed by the relevant specialist locally and unblinding can be performed. It should be noted that trial medication can be withheld for up to 3 months while decisions about subsequent management are confirmed. To discuss individual cases please contact the Asp-PSC team at the CCTS during working hours.

5.5.3 Ischaemic Stroke / Transient Ischaemic Attack (TIA)

The treatment of an ischaemic stroke or TIA frequently involves the use of aspirin and/or other anti-platelet medication. Standard treatment for an ischaemic stroke or TIA should be initiated without delay, and this may include administration of standard doses of anti-platelet therapy, including aspirin, where indicated. It is possible that a trial participant may have taken up to a maximum of 75mg of aspirin earlier that day but, under most circumstances, we expect the risks of not treating an ischaemic stroke or TIA to outweigh the potential risks of a higher cumulative dose of aspirin over one day, however local clinical judgement should be exercised. It is not expected that emergency unblinding will be necessary in the event of an ischaemic stroke or TIA, however if a clinical situation arises where knowledge of the trial treatment allocation would alter the immediate clinical management of a participant, this should be confirmed by the relevant specialist locally and unblinding can be performed. It should be noted that trial medication can be withheld for up to 3 months while decisions about subsequent management are confirmed. To discuss individual cases with the trial's vascular advisors please contact the Asp-PSC team at the Trials Unit during working hours.

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

5.5.4 Surgery

When a surgical intervention is planned, anti-platelet therapy is often withheld temporarily. Trial treatment can be withheld for up to 3 months but please ensure that the Trials Unit is informed. For participants who require emergency surgery, where knowledge of treatment allocation would alter immediate surgical management, this should be confirmed by the surgical team involved locally and unblinding can be performed.

5.6 VISIT SCHEDULE

The table below shows an outline of the proposed study visit schedule.

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

Study Period Screening

→

Time-point → Assessments ↓	Day-0-60	Collection	Week 4 (month 1)	Every 24 Weeks(+ 5 days)	Every 52 weeks (+/-8 weeks)	End of Treatment: 28 days (EOT) after last study dose	Yearly follow up for 5 years post EOT
Randomisation	X						
Informed Consent	X						
Medical History	X		X	X		X	
Demographics	X						
Vital Signs (Body temperature, pulse rate, respiration rate, blood pressure)	X		X ⁶	X		X	
Height	X						
Weight	X			X		X	
Targeted Physical Exam (if clinically indicated)	X		X ⁶	X		X	
Haematology including coagulation ¹	X		X	X			
Biochemistry ¹	X		X	X			
LFTs including AST if locally available ¹	X		X	X			
Tumour markers Ca 19-9, AFP ¹	X			X			
Adverse Events	X		X	X			
Concomitant Medications	X		X	X		X	
QOL Assessments (PSC-PRO, SF- 36, 5D-Itch Scale, SIBDQ, CLDQ- PSC and PRO-2)	X		X	X		X	
Research Urine Sample Collection	X			X			
Research Blood Collection	X			X			
US/Dynamic MRI Liver/Dynamic CT within 6 months of randomisation date	X ²						

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

US liver, MRCP or dynamic MRI/CT within 12 months of randomisation date	X ³						
US/MRI/MRCP/CT at follow up				X ²	X ³		
ERCP/Spyglass Cholangioscopy result collection ⁴	X			X		X	
Colonoscopy	X ⁵				X		
Record of treatment endpoints and survival				X		X	X
Non-invasive assessment of liver fibrosis (serum ELF test)	X						
UKELD Score	X			X			
Stool sample for faecal calprotectin					X		
Fibroscan result collection if available ⁵	X			X			
Collection of aspirin/placebo		X ⁷		X			

1. Only collect bloods if this is standard of care, if not please use last set of bloods for the eCRF
2. Cirrhosis only Patient
3. Patient without Cirrhosis
4. As per standard of care
5. If locally available, use last scan available from 12 months of randomisation date.
6. Patient may have assessment if clinically needed after month one phone call.
7. Participant's Contact Card and Questionnaires for one month visit should also be given.

5.7 FOLLOW-UP

After successful randomisation, patients will attend a collection visit for collection of aspirin/placebo, please ensure the participant contact's card is given and the questionnaires to be completed at month one phone call.

Patients will be called at month one, 4 weeks after the collection visit. The first month's phone call will be to ensure drug tolerability and check no immediate adverse events.

During the month one phone call, the patients will be asked for:

1. Medical History
2. Review of Haematology including FBR and INR test if locally available
3. Review of Biochemistry including UE, LFT, GGT, AST, CRP, Vitamin D, B12, ferritin, TSH and T4 and if locally available
4. Adverse Events
5. Concomitant Medications

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

6. QOL Assessments (PSC-PRO, SF-36, 5D-Itch Scale, SIBDQ, PRO-2 and CLDQ-PSC)

During the one-month phone call patients may be required to attend an in person visit if clinically required to assess further adverse event reporting, this will be decided by a delegated member of staff responsible for patient care. The following assessments will be conducted:

1. Vital Signs (Body temperature, pulse rate, respiration rate, blood pressure-if clinically indicated))
2. Targeted Physical Exam (if clinically indicated)

One-month questionnaires can be administered in the following ways:

- Given to the patient at the collection visit.
- Mailed to the patient's home in time for the one-month phone call.
- Administered over the phone by a designated staff member, who will record the patient's responses on a paper questionnaire. If this method is used, the staff member must note their name, signature, and date on the questionnaire.

After this time point, patients will be seen at 6 monthly intervals (5 months after month 1 phone call, then 6 monthly thereafter) ideally in alignment with routine clinical care during which a research visit will be undertaken.

During each 6-month visit, the patients will be asked to:

1. Documentation of medical history
2. Documentation of vital signs (Body temperature, pulse rate, respiration rate, blood pressure)
3. Weight measurement
4. Targeted Physical Exam (if clinically indicated)
5. Undertake standard of care National Health Service (NHS) blood tests (Full Blood Count (FBC), Urea and Electrolytes (U&E), liver biochemistry, gamma-glutamyl transferase (GGT), aspartate aminotransferase (AST), C-Reactive Protein (CRP), International Normalised Ratio (INR), Vitamin D, Cancer Antigen 19.9 (CA19.9), Alpha-fetoprotein (AFP), ferritin and a TSH and T4). Standard of care faecal calprotectin measurement will be taken yearly.
6. Adverse event documentation
7. Complete questionnaires: Primary Sclerosing Cholangitis -Patient reported outcome (PSC PRO), Short Form-36 (SF-36), 5D-Itch Scale, and Short Inflammatory bowel disease questionnaire (SIBDQ), PRO-2 (Patient Reported Outcome-2) and Chronic Liver Disease Questionnaire (CLDQ)-PSC. Questionnaires can be taken home to do. If Questionnaires are not brought to clinic, site will need to ask patient to complete questionnaires on the day of visit.
8. Documentation of Concomitant Medications
9. Urine and blood samples taken (if patient consented to this).
10. Body mass index (BMI) measurement recorded.

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

11. Patients will also be asked about their alcohol and smoking habits.
12. A fibroscan will also be undertaken if locally available.
13. Documentation of UKELD Score
14. Note will be made of any intervening cancer diagnosis, decompensating liver events or referral for liver transplantation.
15. Note will also be made of any colonoscopy undertaken and also the histological findings from that examination. Note will also be made of any endoscopic intervention/cholecystectomy or new cancer diagnosis made.
16. Note will also be made of any recent imaging studies the patients have undergone, the results available from these studies and multidisciplinary team (MDT) outcomes.
17. Collection of aspirin/placebo tablets
18. Return of the one month questionnaire (if completed at home)

Certain assessments will be required every 12 months following standard of care practice:

1. US/MRI/MRCP/CT at follow up if the patient does not have Cirrhosis
2. Data from colonoscopy
3. Stool sample for faecal calprotectin test

An end of treatment visit will be performed 28 days after last study dose:

4. Documentation of medical history
5. Documentation of vital signs (Body temperature, pulse rate, respiration rate, blood pressure)
6. Weight measurement
7. Targeted Physical Exam (if clinically indicated)
8. Documentation of Concomitant medication
9. QOL Assessments (PSC-PRO, SF-36, 5D-Itch Scale, SIBDQ, CLDQ-PSC and PRO-2)
10. ERCP/Spyglass Cholangioscopy result collection as standard of care.
11. Record of treatment endpoints and survival

5.7.1 Laboratory Evaluations

The blood tests (FBC, UE, LFT, GGT, AST, CRP, INR, Vitamin D, CA19.9, AFP, ferritin and TSH and T4) will be performed locally and the results will be recorded on the eCRF.

The conduct of these analyses will be in accordance to the local NHS laboratory and accredited pathology service. These will be performed if these bloods are standard of care, ELF testing will be conducted at screening and sent to a central laboratory for testing at iQur and results will be sent back to the local site. This will also be used centrally for the randomisation process. Full instructions will be provided in a laboratory manual.

11ml of blood will be taken at screening visit and 43ml of blood will be taken every 6 month follow up for 5 years. 10ml of Urine samples will be taken at screening visit and every 6 months for 5 years. These will be collected in accordance to the separate laboratory manual. These will be stored at participating sites at -80 degrees Celsius and shipped annually on dry ice to the Norwich Biorepository. Biosamples will be made accessible to

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

other research groups via an application process. Patients may choose not to donate research samples but still remain on the trial. A serum sample will be sent to iQur for ELF testing. This is not an optional sample.

- i) **Haematology:** This will include local labs for FBC and INR at standard of care visits.
- ii) **Biochemistry:** This will include U&E, Liver function tests, GGT, aspartate transaminase, CRP, vitamin D, CA19.9, AFP, B12 a ferritin and TSH and T4 at standard of care visits.
- iii) **Research Urine sample:** A urine sample is to be performed in all participants at screening and each study visit in accordance with the laboratory manual. Samples will be collected and stored in accordance with the laboratory manual. Samples will be shipped annually to the Norwich Biorepository.

In all cases, patients will be consented for the **collection and use of their biological samples** and a full chain of custody will be maintained for all samples throughout their lifecycle. The Investigator at each site is responsible for maintaining a record of full traceability of biological samples collected from patients while these are in storage at the site, either until shipment or disposal. Any person(s) responsible for temporarily holding samples, e.g. sub-contracted service provider must keep full traceability of samples from initial receipt of sample to further shipment or disposal (as appropriate).

Norwich Research Park Biorepository will keep overall oversight of research Biosamples the entire lifecycle through internal procedures and monitoring of study sites. iQur will keep oversight of the ELF testing sample.

Other central laboratories may be used for ELF testing including South West Pathology, Cambridge University Hospital pathology OR Black Country Pathology.

5.8 INCIDENTAL FINDINGS

Any incidental findings will be identified and reviewed and acted upon by the local PI who will determine if they should be handled and investigated by the local clinical team, a different secondary NHS team or passed to the primary care team.

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

6 TREATMENTS

6.1 TREATMENT ARMS

Within this study an oral formulation of enteric coated aspirin will be compared to placebo (manufactured so that it looks identical) by the designated provider. IMP will be dispensed in bottles. The medication will be distributed to participating sites via processes managed by ICTU during the screening period with the patient being randomised ahead of collection visit. The IMP will be manufactured under Good Manufacturing Practice by The Royal Free Hospital Pharmacy. Currently the legal status of aspirin is licensed for approval in the UK. For the purposes of this trial the Summary of product characteristics (SmPC) will be used for regulatory approval.

A list of side effects can be found in appendix 9 in SmPC.

6.2 LABELLING AND PACKAGING

The IMP will be packaged and labelled in bottles ahead of distribution of participating sites. The label will contain:

- Manufacturing Import Authorisation (MIA) (IMP)
- For Clinical Trials Use Only
- Aspirin EC 75mg/Placebo Tablets
- Store at Room Temperature, under 25°C
- 92 Tablets
- Asp-PSC Trial
- IRAS Number:
- Sponsor:
- Principal Investigator:
- Site Name/Number:
- Subject ID/Randomisation ID:
- Batch Number:
- Expiry:
- Kit Number:
- For Oral Use only
- Internal Label code:
- Date dispensed:
- Keep out of reach and sight of children

6.3 STORAGE AND DISPENSING

Information regarding storage and dispensing will be found in the IMP manual provided to sites.

6.4 DOSAGE, DURATION AND COMPLIANCE

The IMP will be taken once a day in the morning or evening with a glass of water. The IMP will be taken orally. All IMP compliance will be recorded daily in a patient diary. The

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

compliance of the drug will be checked at each 6-monthly visit, where the number of tablets left in the boxes will be manually counted and recorded in the eCRF. Post study (5 years) the IMP will be withdrawn.

6.5 ACCOUNTABILITY

The IMP is only to be prescribed by the PI or delegated investigator and dispensed by delegated members of the research team at the participating site for participants within the trial as specified in this protocol. ICTU will facilitate IMP shipment to participating sites via the distributor. A full accountability trail of the trial medication will be maintained from receipt at the participating site until destruction.

6.6 DRUG INTERACTIONS

See appendix 9 for drug interactions.

6.6.1 Contraindicated Medications

Patients should not take any of the following contraindicated medications whilst taking aspirin/placebo:

- Concurrent use of other antiplatelet or anticoagulant agents
- Aspirin
- Methotrexate at doses of more than 15mg/week
- Uricosuric agents such as probenecid and sulfinpyrazone
- NSAIDs-chronic daily long term usage (>3 weeks)

Patients should not enrol on another CTIMP study whilst taking aspirin or placebo on trial.

6.6.2 Need to prescribe concurrent oral corticosteroids

If patient who has been enrolled into this study needs corticosteroids prescribing, they should be assumed to be taking aspirin as their IMP. All patients therefore prescribed oral corticosteroids should be co-prescribed a proton pump inhibitor (or Histamine 2 Receptor blocker if intolerant) as prophylaxis against Gastrointestinal bleeding for the duration that corticosteroid is prescribed.

6.7 DOSE MODIFICATIONS FOR ADVERSE EVENTS

This is not applicable in this study.

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

6.8 OVERDOSE OF IMP

Although considerable inter-individual variations are involved, it can be considered that the toxic dose of aspirin is about 200 mg/kg in adults. The lethal dose of acetylsalicylic acid is 25-30 grams. Salicylate poisoning is usually associated with plasma concentrations >350 mg/L (2.5 mmol/L). Plasma concentrations above 500 mg/l in adults generally cause severe toxicity. Most adult deaths occur in patients whose concentrations exceed 700 mg/L (5.1 mmol/L). Single doses less than 100 mg/kg are unlikely to cause serious poisoning.

Plasma salicylate concentrations should be measured urgently for patients who are thought to have ingested more than 125 mg/kg of aspirin. The sample should be taken at least 2 hours (symptomatic patients) or 4 hours (asymptomatic patients) after ingestion, since it may take several hours for peak plasma concentrations to occur **and up to 12 hours for enteric-coated preparations**. A repeat sample should be taken in ALL symptomatic patients and those with concentrations greater than 500 mg/L after a further 2 hours because of the possibility of continuing absorption. Under these circumstances, measurements should be repeated every 3 hours until concentrations are falling.

6.8.1 Symptoms

Common features include vomiting, dehydration, tinnitus, vertigo, deafness, sweating, warm extremities with bounding pulses, increased respiratory rate and hyperventilation. Some degree of acid-base disturbance is present in most cases.

A mixed respiratory alkalosis and metabolic acidosis with normal or high arterial pH (normal or reduced hydrogen ion concentration) is usual in adults. Acidosis may increase salicylate transfer across the blood brain barrier.

Uncommon features include haematemesis, hyperpyrexia, hypoglycaemia, hypokalaemia, thrombocytopaenia, increased INR/PTR, intravascular coagulation, renal failure and non-cardiac pulmonary oedema.

Other symptoms may include: headache, nausea, or abdominal pain.

Central nervous system features including confusion, restlessness, hallucinations, disorientation, coma, cardiovascular collapse, respiratory arrest and convulsions are less common in adults than in children.

If a toxic dose has been ingested, hospital admission is required.

Activated charcoal should be given if a patient presents within one hour of ingestion of more than 250 mg/kg. The plasma salicylate concentration should be measured, although the severity of poisoning cannot be determined from this alone and the clinical and biochemical features must be considered. Elimination is increased by urinary alkalinisation, which is achieved by the administration of 1.26% sodium bicarbonate. The urine pH should be monitored. Correct metabolic acidosis with intravenous 8.4% sodium bicarbonate (first check serum potassium). Forced diuresis should not be used since it does not enhance salicylate excretion and may cause pulmonary oedema.

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

Haemodialysis is the treatment of choice for severe poisoning and should be considered in patients with plasma salicylate concentrations >700 mg/L (5.1 mmol/L), or lower concentrations associated with severe clinical or metabolic features. Patients over 70 have increased risk of salicylate toxicity and may require dialysis at an earlier stage.

Other symptoms should be treated symptomatically.

6.9 PERMANENT DISCONTINUATION OF STUDY TREATMENT

Participants may discontinue study treatment for the following reasons:

- At the request of the participant
- Adverse Event/ Serious Adverse Event
- Allergic reaction to IMP
- If the investigator considers that a participant's health will be compromised due to adverse events or concomitant illness that develop after entering the study.
- In the event they are prescribed another anti-platelet agent or anti-coagulant

The primary reason for discontinuation must be recorded in the eCRF and the patient's medical records. If a participant chooses to discontinue their trial treatment, they should continue to be followed up as according to the schedule defined in the protocol. Every effort should be made to encourage participants not to withdraw from the study.

6.9.1 Withdrawal from Study

Withdrawal from the study refers to discontinuation of study treatment and study procedures and can occur for the following reasons:

- Participant decision
- Investigator Decision
- Loss to follow-up
- Death

Data will still be collected for participants for follow up unless they withdraw from further data collection.

If a participant dies whilst participating in the study, a 'Statement of Death' eCRF must be completed. The following information will be collected: date of death, cause of death and if the death is related to disease.

Reaching the following endpoints will withdraw the patient from the study, but the patient will then enter yearly follow up:

Endpoint	Withdraw patient yes/no
Variceal Bleeding	Yes
Ascites	Yes
Hepatic encephalopathy	No
Development of cirrhosis	No

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

Acute cholangitis	No
IBD flares	No
Colonic Resection	Withdraw patient if performed for cancer or dysplasia if performed for severe IBD then do not withdraw the patient.

6.9.2 Procedures for Withdrawal from Study

In the event of a patient withdrawing from the study, no further visits or follow up will be conducted. If the patient only withdraws from taking IMP, patient can enter long term follow up, see section 6.9.1. The primary reason must be recorded in the eCRF and the patient's medical records. Any data and samples already collected will be retained and analysed.

Where a patient has been lost to follow-up, this should be recorded on the eCRF with their last known contact date and survival status at the time. If appropriate, patients who withdraw will be replaced.

6.9.3 Temporary Discontinuation of IMP in the study

Patients will be allowed to discontinue their IMP for up to a 3-month period if required for clinical reasons, for example the need for surgery or if they become pregnant and wish to discontinue the IMP for the first trimester, and then continue for a total of five years follow-up.

7 PHARMACOVIGILANCE

7.1 ADVERSE EVENT (AE)

An AE is any untoward medical occurrence in a patient or clinical trial participant administered a medicinal product and which does not necessarily have a causal relationship with this treatment. An AE can therefore be any unfavourable and unintended sign (including an abnormal laboratory finding), symptom, or disease temporally associated with the use of the trial medication, whether or not considered related to the IMP.

7.2 ADVERSE REACTION (AR)

All untoward and unintended responses to an IMP related to any dose administered. All AEs judged by either the reporting investigator or the sponsor as having reasonable causal relationship to a medicinal product qualify as adverse reactions (ARs). The expression of reasonable causal relationship means to convey in general that there is evidence or argument to suggest a causal relationship.

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

7.3 UNEXPECTED ADVERSE REACTION

An AR, the nature or severity of which is not consistent with the applicable product information as set out in the Reference Safety Information (RSI) i.e. summary of product characteristics (SmPC).

When the outcome of the adverse reaction is not consistent with the applicable product information this adverse reaction should be considered as unexpected. **Side effects documented in the RSI section of the SmPC which occur in a more severe form than anticipated are also considered to be unexpected.**

Expectedness assessment will be performed by the Sponsor or person delegated by the Sponsor to assess expectedness.

For this study this will be the local PI who makes a determination of this.

7.4 CAUSALITY

The assignment of causality for adverse events should be made by the investigator responsible for the care of the participant using the definitions in the table below.

If any doubt about the causality exists, the local investigator should inform ICTU who will notify the Chief Investigator. The pharmaceutical companies and/or other clinicians may be asked to advise in some cases.

In the case of discrepant views on causality between the investigator and others, all parties will discuss the case. In the event that no agreement is made, the Medicines and Healthcare Products Regulatory Agency (MHRA) will be informed of both points of view. For the purpose of causality, the following definitions will be used.

- Unrelated: No evidence of any causal relationship
- Unlikely: There is little evidence to suggest there is a causal relationship (e.g., the event did not occur within a reasonable time after administration of the trial medication). There is another reasonable explanation for the event (e.g. the patient’s clinical condition, other concomitant treatment).
- Possible: There is some evidence to suggest a causal relationship (e.g., because the event occurs within a reasonable time after administration of the trial medication). However, the influence of other factors may have contributed to the event (e.g., the patient’s clinical condition, other concomitant treatments).
- Probable: There is evidence to suggest a causal relationship and the influence of other factors is unlikely.
- Definite: There is clear evidence to suggest a causal relationship and other possible contributing factors can be ruled out.

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

7.5 SEVERITY OF ADVERSE EVENTS

Severity of AEs will be assessed using the grading scales found in the National Cancer Institute CTCAE version 5.0 (27th November 2017), by attributing the most relevant CTCAE term, Grade 1 = Mild, Grade 2 = Moderate, Grade 3 = Severe, Grade 4 = Life Threatening and Grade 5 = Death. Where it is not possible to attribute a specific CTCAE term, a term may be attributed by the investigator and assessed according to the introduction under Grades in CTAE version 5.0 i.e., grade 1=mild; grade 2=moderate; grade 3=severe; grade 4=life-threatening; grade 5= fatal.

CTCAE version 5.0 is accessible online here:

https://ctep.cancer.gov/protocoldevelopment/electronic_applications/ctc.htm

7.6 ADVERSE EVENT RECORDING

All AEs should be recorded in the patient's medical notes and entered into the eCRF within 10 days.

Adverse Event (AE)	Any untoward medical occurrence in a patient or clinical trial participant administered a medicinal product and which does not necessarily have a causal relationship with this product.
Adverse Reaction (AR)	Any untoward and unintended response to an investigational medicinal product related to any dose administered/trial treatment
Unexpected Adverse Reaction (UAR)	An adverse reaction, the nature or severity of which is not consistent with the applicable product information (e.g. Investigator's Brochure for an unauthorised product or summary of product characteristics (SPC) for an authorised product or treatment
Serious Adverse Event (SAE) or Serious Adverse Reaction (SAR)	Any AE or AR that at any dose: <ul style="list-style-type: none"> • results in death • is life threatening* • requires hospitalisation or prolongs existing hospitalisation** • results in persistent or significant disability or incapacity • is a congenital anomaly or birth defect • or is another important medical condition***
<p>* The term life threatening here refers to an event in which the patient is at risk of death at the time of the event; it does not refer to an event that might hypothetically cause death if it was more severe (e.g., a silent myocardial infarction)</p> <p>** Hospitalisation is defined as an in-patient admission, regardless of length of stay, even if the hospitalisation is a precautionary measure for continued observation.</p>	

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

Hospitalisation for pre-existing conditions (including elective procedures that have not worsened) do not constitute an SAE

*** Medical judgement should be exercised in deciding whether an AE or AR is serious in other situations. Important AEs or ARs that may not be immediately life threatening or result in death or hospitalisation, but may seriously jeopardise the participant by requiring intervention to prevent one of the other outcomes listed in the table (e.g., a secondary malignancy, an allergic bronchospasm requiring intensive emergency treatment, seizures or blood dyscrasias that do not require hospitalisation, or development of drug dependency).

If the event is classified as 'serious' then an SAE form must be completed within the eCRF within 24 hours of the site becoming aware.

Severity or grading of Adverse Events

The severity of all AEs in this trial should be graded using the adverse event gradings in v5.0 CTCAE criteria.

For the purposes of the study, AEs will be followed up according to local practice until the event has stabilised or resolved, or the Follow-up Visit, whichever is sooner. SAEs will be recorded up to 28 days after the last study dose.

7.7 ABNORMAL LABORATORY TEST RESULTS

All clinically important abnormal laboratory test results occurring during the study will be recorded as adverse events. The clinically important abnormal laboratory tests will be repeated at appropriate intervals until they return either to b or to a level deemed acceptable by the investigator and the clinical monitor, or until a diagnosis that explains them is made.

7.8 SERIOUS ADVERSE EVENTS (SAE)

7.8.1 Definition of SAE

An SAE is defined as any event that

- Results in death;
- Is life-threatening*;
- Requires hospitalisation or prolongation of existing inpatient's hospitalisation**;
- Results in persistent or significant disability or incapacity;
- Is a congenital abnormality or birth defect;

* "Life-threatening" in the definition of "serious" refers to an event in which the participant was at risk of death at the time of the event; it does not refer to an event which hypothetically might have caused death if it were more severe.

** "Hospitalisation" means any unexpected admission to a hospital department. It does not usually apply to scheduled admissions that were planned before study inclusion or visits to casualty (without admission).

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

Medical judgement should be exercised in deciding whether an adverse event/reaction is serious in other situations. Important adverse events/reactions that are not immediately life-threatening, or do not result in death or hospitalisation but may jeopardise a subject, or may require intervention to prevent one of the other outcomes listed in the definition above should also be considered serious.

7.9 REPORTING OF SAEs

Rapid reporting of all SAEs i.e., within 24 hours of the site becoming aware, occurring during the study must be performed as detailed in the study-specific pharmacovigilance manual.

SAE will be reported from time of randomisation until 28 days after last study dose.

Active monitoring of participants after the end of the trial is not required but if the investigator becomes aware of safety information that appears to be drug related, involving a participant who participated in the study, even after an individual participant has completed the study, this should be reported to the Sponsor.

All SAEs will be reviewed by the Chief Investigator or a designated medically qualified representative to confirm expectedness and causality.

Reporting of SAEs and review by the CI will be via the trial data collection system (eCRF).

7.10 DEFINITION OF A SERIOUS ADVERSE REACTION (SAR)

A SAR is defined as a SAE that is judged to be related to any dose of study drug administered to the participant.

7.10.1 Definition of Suspected Unexpected Serious Adverse Reaction (SUSAR)

Any SAR that is NOT consistent with the applicable product information as set out in the Reference Safety Information (RSI) section of the Investigator Brochure (IB) or Summary of Product Characteristics (SPC).

7.10.2 Reporting of SUSARs

SUSARs should be notified to the appropriate **regulatory authority, the relevant REC and the Sponsor in accordance with regulatory requirements**. SUSARs which are fatal or life-threatening will be reported not later than seven days after alerting the sponsor to the reaction. Any additional relevant information will be sent within eight days of the report.

A SUSAR which is not fatal or life-threatening will be reported within 15 days of first knowledge by the sponsor. The sponsor will inform all investigators about SUSARs occurring on the study. This will be by means of email to all study investigators. Follow up of participants who have experienced a SUSAR should continue until recovery is complete or the condition has stabilised. SUSAR reports should be unblinded prior to submission.

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

7.11 DEVELOPMENTAL SAFETY UPDATE REPORTS / ANNUAL SAFETY REPORTS

Developmental Safety Update Reports (DSUR) / Annual Safety reports will be submitted to the Sponsor, the Ethics Committee and Regulatory Authority in accordance with local / national regulatory requirements.

7.12 PREGNANCY

At the IMP dose used there is no need to avoid or use additional contraception for this study. There are convincing data to show that low-dose aspirin (150 or 75mg daily) protects against severe pre-eclampsia and foetal growth restriction if commenced before 16 weeks' gestation (27)(28). Furthermore, a meta-analysis of eight RCTs that gave low-dose aspirin before 14 weeks to 7564 participants compared to 7670 participants that served as controls reported no significant difference in the rate of congenital malformations (29). Thus participants planning pregnancy, or those who become pregnant will not be asked to stop treatment, although they may stop taking the study drug before 11 weeks if they elect to do this. Pregnancy outcome data relating to maternal disease severity and maternal and fetal outcomes will be collected from trial participants. Participants can also continue to breastfeed if they wish to do so.

If a patient on trial becomes pregnant, they should be referred to the Obstetric Team for enhanced antenatal care in line with UK clinical practice for women with chronic liver disease.

7.13 REPORTING URGENT SAFETY MEASURES

If any urgent safety measures are taken the CI/Sponsor shall immediately and in any event no later than 3 days from the date the measures are taken, give written notice to the MHRA and the relevant REC of the measures taken and the circumstances giving rise to those measures.

8 STATISTICAL ANALYSES

8.1 DESIGN

This study will be a multicentre, double-blind, randomised placebo-controlled trial. The study population will include patients with PSC who have a concurrent diagnosis of IBD and are at least 12 months post diagnosis of PSC. Patients will be randomised in a ratio of 2:1 Aspirin vs Placebo. Accrual will be over 5 years, with a minimum participant follow up of 5 years. Included in this study is an embedded feasibility study for 12 months from the start of recruitment to establish recruitment, acceptability and compliance rates.

Randomisation will be stratified by the following factors: 1) degree of fibrosis (binary, as determined by ELF score >9.8, OR Fibroscan score (if ELF test cannot be performed) of 9.6 kPa), PSC disease duration (as determined by 1 - <5 , ≥5 - <10, ≥10) and concurrent use of aminosalicylates (yes or no).

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

8.2 SAMPLE SIZE AND POWER CONSIDERATIONS

774 patients need to be recruited to the trial (2:1 randomisation) to allow us to detect, with 80% power (alpha 0.05), a HR of 0.6, equivalent to a difference between groups of 6% in the proportion of patients experiencing any of the components of the primary outcome, namely a new cancer diagnosis, or requiring liver transplant or any cause mortality. The intervention:placebo ratio is 2:1, meaning we will need to recruit 516 participants in the intervention group and 258 in the placebo group. This is based on an estimated event rate in the control group of 15% at 5 years (2), 5 years of accrual and 5 years of minimum follow up. To allow for 20% loss to follow up, we have determined we will need to recruit 968 patients.

8.3 PLANNED RECRUITMENT RATE

See section 4.2 for recruitment table.

8.4 STATISTICAL ANALYSIS

A Statistical Analysis Plan (SAP) will be written and approved before data lock. This document will outline all analyses for all the outcomes listed in the Protocol.

8.5 ANALYSIS OF PRIMARY AND SECONDARY OUTCOMES

Summaries of continuous variables will be presented as mean and standard deviations if normally distributed, and as medians and inter-quartile ranges for skewed data. Categorical variables will be presented as frequencies and percentages in the form of summary tables. Time to event endpoints will be graphically displayed using Kaplan-Meier curves comparing treatment groups. Median time to event and 2-sided 95% confidence intervals will be reported. All statistical tests will be two-tailed with a 5% significance level. A Cox proportional hazard model will be used to estimate the HR between the two groups. The model will be adjusted by the randomisation stratification factors and centre. If the assumption for the Cox model are not met we will use an alternative model.

8.6 PRIMARY EFFICACY OUTCOME ANALYSIS

The composite primary endpoint is defined as occurrence of any of the following components:

1. Hepatobiliary cancer (including gallbladder cancer/high grade dysplasia, pancreatic cancer, cholangiocarcinoma [CCA] or HCC) or colorectal cancer/high grade dysplasia
2. Liver transplantation
3. All-cause mortality

Using the Kaplan-Meier method, curves for the combined event free survival (where events are cancer/high grade dysplasia need of liver transplant and overall mortality will be

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

presented for each treatment arm, with time from randomisation until the first failure event. The primary inferential comparison between treatment groups will estimate the HR with 95% CI. The HR will be estimated using a Cox proportional hazards model adjusted by the stratification factors, if the proportional hazard assumptions are met. If the proportional hazard assumptions are not met alternative methods will be used to estimate the difference between groups.

8.7 SECONDARY OUTCOMES' ANALYSIS

Secondary endpoints include GI bleeding, progressive liver disease, IBD flares including the need for colectomy, referral for liver transplantation, acute cholangitis and quality of life.

This will be done by using the chi-squared test and logistic regression adjusted by stratification factors and other pre-defined factors. Quality of life data will be analysed using a mixed regression model, adjusted by stratification factors and other predefined factors, to take into account the repeated measures in time.

A Statistical Analysis Plan (SAP) will be written ahead of looking at the data, and will be signed off by the research team and shared with the DMEC and TSC.

8.8 MISSING DATA

Appropriate methods of imputation will be used to deal with missing data, if the missing data are more than 20%. Details of the type of analysis will be included in the SAP.

9 REGULATORY, ETHICAL AND LEGAL ISSUE

9.1 DECLARATION OF HELSINKI

The investigator will ensure that this study is conducted in full conformity with the 1964 Declaration of Helsinki and all subsequent revisions.

9.2 GOOD CLINICAL PRACTICE

The study will be conducted in accordance with the guidelines laid down by the International Conference on Harmonisation for Good Clinical Practice (ICH GCP E6 guidelines).

9.3 RESEARCH ETHICS COMMITTEE (REC) APPROVAL

9.3.1 Initial Approval

Prior to the shipment of IMP and the enrolment of participants, the REC must provide written approval of the conduct of the study at named sites, the protocol and any amendments, the Participant Information Sheet and Consent Form, any other written information that will be provided to the participants, any advertisements that will be used and details of any participant compensation.

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

9.3.2 Approval of Amendments

Proposed amendments to the protocol and aforementioned documents must be submitted to the REC for approval. Amendments requiring REC approval may be implemented only after a copy of the REC's approval letter has been obtained.

Amendments that are intended to eliminate an apparent immediate hazard to participants may be implemented prior to receiving Sponsor or REC approval. However, in this case, approval must be obtained as soon as possible after implementation.

Amendments to the Protocol and other documents (e.g., changes to eligibility criteria, outcomes, sample size calculations, analyses) will be agreed by the TMG. Such amendments will be forwarded to the Sponsor for confirmation as to whether it is either substantial or non-substantial and will then be submitted to the Health Research Authority or Ethics Committee for categorisation and approval.

Once the amendment has been categorised it will be sent to relevant sites for consideration in accordance with standard HRA processes and timescales. Amendments must not be implemented until HRA approval is received and sites have either confirmed acceptance or, no objection has been received within the defined timescale. Notification will be sent by CTS to trial personnel to confirm when an amendment can be implemented.

9.3.3 End of Trial Notification

The REC will be informed about the end of the trial, within the required timelines.

The end of trial notification will be submitted within 90 days of the end of trial definition being met. In the event of a premature halt of the trial, the timeframe is 15 days, and the reasons should be clearly explained in the notification.

9.4 REGULATORY AUTHORITY APPROVAL

The study will be performed in compliance with United Kingdom regulatory requirements. Clinical Trial Authorisation from the appropriate Regulatory authorities must be sought prior to the start of the study. In addition, the Regulatory Authority must approve amendments prior to their implementation (as instructed by the Sponsor), receive SUSAR reports and annual safety updates and be notified of the end of the trial. This study has Clinical Trials Authorisation from the UK Competent Authority; MHRA. **Reference:** CTA 19174/0447/001-0001

9.5 HRA APPROVAL

HRA approval will be obtained prior to starting the study. Each participating site will confirm capacity and capability prior to commencing. The HRA and all participating sites also need to be notified of all protocol amendments to assess whether the amendment affects the institutional approval for each site.

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

9.6 NON-COMPLIANCE AND SERIOUS BREACHES

All protocol deviations and protocol violations will be reported via the eCRF and reviewed by the Chief Investigator and reported to the ICTU Quality Assurance (QA) Manager on a monthly basis. Protocol violations will be reported to the Sponsor. An assessment of whether the protocol deviation/violation constitutes a serious breach will be made.

A serious breach is defined as:

- A breach of the conditions and principles of GCP in connection with a trial or the trial protocol, which is likely to affect to a significant degree:
- The safety or physical or mental integrity of the UK trial participants; or
- The overall scientific value of the trial

The Sponsor will be notified within 24 hours of identifying a likely Serious Breach. If a decision is made that the incident constitutes a Serious Breach, this will be reported to the MHRA and REC within 7 days of becoming aware of the serious breach.

9.7 INSURANCE AND INDEMNITY AND SPONSOR

For the purposes of this trial the sponsor will be:

Imperial College London
Research Governance and Integrity team
Room 221, Medical School Building
St Marys Campus, Norfolk Place
London W2 1PG
United Kingdom

Please note that Imperial College London holds negligent harm and non-negligent harm insurance policies which apply to this study.

Delegated responsibilities will be assigned to the NHS trusts taking part in the trial.

9.8 TRIAL REGISTRATION

The study will be registered on the following trial databases in accordance with requirements of the International Committee of Medical Journal Editors (ICMJE) regulations:

- Cancer Research UK
- ISRCTN

9.9 INFORMED CONSENT

Patients will be provided with a Patient Information Sheet (PIS) and given time to read it fully. Following a discussion with a medical qualified investigator, any questions will be satisfactorily answered and if the participant is willing to participate, written informed consent will be obtained. During the consent process it will be made completely and

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

unambiguously clear that the participant is free to refuse to participate in all or any aspect of the trial, at any time and for any reason, without incurring any penalty or affecting their treatment.

Consent will be re-sought if significant information becomes available that affects the participant's consent in any way. This will be documented in a revision to the patient information sheet and the participant will be asked to sign an updated consent form. These will be approved by the ethics committee prior to their use.

Informed consent will be obtained either in person by a member of the research team who is a medically qualified investigator either the PI or sub-investigator, at the affiliated NHS site using a paper consent form.

9.10 CONTACT WITH GENERAL PRACTITIONER

Once the patient has given full voluntary consent to participate in the trial, the local PI or delegated member of staff will send a GP letter to the patient's GP. This will inform the GP that the participant is taking part in the study, and information to this effect is included in the Participant Information Sheet and Informed Consent. A copy of the GP letter should be filed in the Investigator Site File.

9.11 PARTICIPANT CONFIDENTIALITY

The investigator must ensure that the participant's confidentiality is maintained. On the eCRF or other documents submitted to the Sponsors, participants will be identified by a participant ID number only. Documents that are not submitted to the Sponsor (e.g., signed informed consent form) should be kept in a strictly confidential file by the investigator.

The investigator shall permit direct access to participants' records and source documents for the purposes of monitoring, auditing, or inspection by the Sponsor, authorised representatives of the Sponsor, NHS, Regulatory Authorities and RECs.

9.12 DATA PROTECTION AND PARTICIPANT CONFIDENTIALITY

The investigators and study site staff will comply with the requirements of the Data Protection Act 2018 concerning the collection, storage, processing and disclosure of personal information and will uphold the Act's core principles.

9.13 END OF TRIAL

For the purposes of this clinical trial, the end of the trial will be when the last participant has completed 5 years follow up since the date of randomisation and the final end of trial visit.

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

9.14 STUDY DOCUMENTATION AND DATA STORAGE

The investigator must retain essential documents until notified by the Sponsor, and for at least ten years after study completion. Participant files and other source data (including copies of protocols, original reports of test results, IMP dispensing logs, correspondence, records of informed consent, and other documents pertaining to the conduct of the study) must be retained. Documents should be stored in such a way that they can be accessed/data retrieved at a later date. Consideration should be given to security and environmental risks.

No study document will be destroyed without prior written agreement between the Sponsor and the investigator. Should the investigator wish to assign the study records to another party or move them to another location, written agreement must be obtained from the Sponsor.

10 DATA MANAGEMENT

10.1 SOURCE DATA

The following source data will be recorded in this study:

- Medical records
- Laboratory reports
- Diaries and questionnaires
- IMP documentation
- Radiological imaging and reports
- Colonoscopy reports and histology reports
- Scan reports
- ELF Results
- Questionnaires

Please note that all of these will need to be recorded on the eCRF.

Copies of standard of care scans completed from randomisation to the end of trial will be collected and sent to Infinitt Europe GMBH for storage.

10.2 LANGUAGE

eCRFs will be in English. Generic names for concomitant medications should be recorded in the eCRF wherever possible. All written material to be used by participants must use vocabulary that is clearly understood and be in the language appropriate for the study site.

10.3 DATABASE

Data will be captured using the electronic Case Report Form (eCRF). Data will be entered under the study ID onto the eCRF held on servers based at ICTU. Access to the database

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

will be via unique, individually assigned (i.e. not generic) usernames and passwords, and only accessible to members of the Asp-PSC trial team, and external regulators if requested. The servers are protected by firewalls and are patched and maintained according to best practice. The physical location of the servers is protected physically and environmentally in accordance with ICTU SOPs.

The database software provides a number of features to help maintain data quality, including maintaining an audit trail, allowing custom validations on all data, allowing users to raise data query requests, and search facilities to identify validation failure/ missing data.

After completion of the trial the database will be retained on the servers of ICTU for on-going analysis of secondary outcomes.

The identification, screening and enrolment logs, linking participant identifiable data to the study ID, will be held locally by the trial site. This will either be held in written form in a locked filing cabinet or electronically in password protected form on hospital computers. After completion of the trial the identification, screening and enrolment logs will be stored securely by the sites for 10 years unless otherwise advised by ICTU.

10.4 DATA COLLECTION

Full details of data collection will be documented in a data management plan approved prior to study opening. Details of procedures for eCRF completion will be provided in a study manual.

10.5 ARCHIVING

All trial documentation, including that held at participating sites and ICTU, will be archived for a minimum of 10 years following the end of the study.

11 STUDY MANAGEMENT STRUCTURE

11.1 TRIAL STEERING COMMITTEE

An independent Trial Steering Committee (TSC) will be convened including as a minimum an independent Chair, independent clinician, independent statistician and two lay members. The role of the TSC is to provide overall supervision of trial conduct and progress. Details of membership, responsibilities and frequency of meetings will be defined in a separate Charter.

11.2 TRIAL MANAGEMENT GROUP

A Trial Management Group (TMG) will be convened including the Chief Investigators, co-investigators and key collaborators, trial statistician/s, ICTU Study Manager, ICTU Operations Manager and the PPIE leads. The TMG will be responsible for day-to-day conduct of the trial and operational issues. Details of membership, responsibilities and frequency of meetings will be defined in separate terms of Reference.

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

11.3 INDEPENDENT DATA MONITORING COMMITTEE

An Independent Data Monitoring Committee (IDMC) will be convened including as a minimum an independent Chair, independent clinician and independent statistician. The role of the IDMC is to monitor the efficacy and safety of study data and report its findings to the TSC. Details of membership, responsibilities and frequency of meetings will be defined in a separate Charter.

11.4 EARLY DISCONTINUATION OF THE STUDY

It is possible that the study will require early discontinuation e.g., on IDMC recommendation to the TSC. Were this to occur then the relevant actions would be completed by ICTU, to include management of patients on active study treatment and informing the Sponsor, REC and MHRA.

11.5 RISK ASSESSMENT

A study-specific risk assessment will be performed prior to the start of the study to assign a risk category of 'low', 'medium' or 'high' to the trial. Risk assessment will be carried out by the ICTU QA Manager in collaboration with the ICTU Study Manager and the result will be used to guide the monitoring plan. The risk assessment will consider all aspects of the study and will be updated as required during the course of the study.

11.6 MONITORING

The study will be monitored periodically by trial monitors to assess the progress of the study, verify adherence to the protocol, ICH GCP E6 guidelines and other national/international requirements and to review the completeness, accuracy and consistency of the data.

11.6.1 Central Monitoring at ICTU

Trial staff will review eCRF data for errors and missing key data points. The trial database will also be programmed to generate reports on errors and error rates. Essential trial issues, events and outputs, including defined key data points, will be detailed in the Asp-PSC trial Data Management Plan.

11.6.2 On-site Monitoring

The frequency, type and intensity of routine and triggered on-site monitoring will be detailed in the Asp-PSC Monitoring Plan. The Monitoring Plan will also detail the procedures for review and sign-off of monitoring reports. In the event of a request for a trial site inspection by any regulatory authority, ICTU must be notified as soon as possible.

11.6.2.1 Direct access to participant records

Participating investigators must agree to allow trial related monitoring, including audits, REC review and regulatory inspections, by providing access to source data and other trial

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

related documentation as required. Participant consent for this must be obtained as part of the informed consent process for the trial.

11.7 QUALITY CONTROL AND QUALITY ASSURANCE

Quality Control will be performed according to ICTU internal procedures. The study may be audited by a Quality Assurance representative of the Sponsor and/or ICTU. All necessary data and documents will be made available for inspection. The study may be participant to inspection and audit by regulatory bodies to ensure adherence to GCP and the NHS Research Governance Framework for Health and Social Care (2nd Edition).

11.8 PEER REVIEW

This protocol was generated from an original grant application which was peer reviewed by CRUK, ahead of the decision to grant funding, by three experts in the field of PSC and CCA. The protocol was also endorsed and reviewed by all members of the TMG.

11.9 PUBLIC INVOLVEMENT

11.9.1 Patient and Public Involvement and Engagement (PPIE)

The PPIE for this study has been by undertaken by Martine Walmsley (PSC patient/ PSC Support); and Helen Morement, Founder and CEO of AMMF – The Cholangiocarcinoma Charity. PSC Support and AMMF are national charities, helping people affected by PSC and CCA through education, support, advocacy and research. Martine and Helen are the PPIE leads for the study. They have been extensively involved in planning this proposal from the conception of the idea. PSC Support highlighted that one of the most difficult aspects of living with PSC for patients is the emotional impact, particularly uncertainty about the future and that reducing cancer risk was very important to people affected by PSC.

The PPIE members have extensive experience: working directly with patients at risk of and living with PSC cancers, running research focus groups for patients and loved ones, working with Healthcare/Research teams including developing and supporting novel research proposals and study protocols, collating feedback from and disseminating research findings to patients, and helping to develop national clinical practice guidelines (including for PSC, CCA and liver transplantation).

They have been extensively involved in developing this proposal from conception so that we could draw on experiences and views of people affected by the risk of PSC-related cancers when planning the trial design. The PPIE leads and affected patients met monthly over two years to understand patient priorities (including the realities of cancer risk and the importance of reducing that risk from a patient perspective), appetite for taking part in longer studies, developing the trial design to ensure it was acceptable to patients, and understanding any concerns.

The PSC Support charity carried out a national survey of almost 200 PSC patients to gauge interest and availability for this trial (<https://www.pscsupport.org.uk/surveys/aspirin-to-prevent-cancer-in-psc>). The survey found that the majority of patients:

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

- i) believe it is 'extremely important' to reduce cancer risk for people with PSC (90%)
- ii) would be interested in taking part in a 5-year study (86%), and
- iii) are not already taking aspirin (87%)

The PPIE members have been integral to our planning and advised on:

- Randomisation ratio (to give participants a higher chance of receiving active drug than placebo)
- Recruitment strategy (for example, what aspects of the design would be appealing or a barrier to participation; appetite for participation in trials)
- Inclusion and exclusion criteria
- Patient Reported Outcome Measures (which tools were preferred by PSC patients following previous focus group work by PSC Support and their willingness to regularly complete multiple questionnaires)
- Relevant outcomes (including utilising this opportunity to collect longitudinal data from PSC patients for future biomarker discovery as there are currently no convincing surrogate endpoints for clinical trials).

11.9.2 Patient & Public Involvement & Engagement (PPIE) Lead

The ICTU Study Manager will act as PPIE Lead along with Martine Walmsley and Helen Morement. Tailored training and support will be given to the leads by the ICTU PPIE Coordinator. The PPIE Leads will undertake a number of activities including: developing role descriptions with PPIE Advisory Group; supporting PPIE training needs; acting as PPIE liaison; ensuring involvement is aligned to UK Standards for Public Involvement; providing impact feedback on activities; planning meetings and keeping a PPIE impact log.

11.9.3 PPIE Collaboration Plan

In addition to our two PPIE Co-Investigators who are embedded as part of the main Investigator Team and Trial Management Group (TMG) for the life of the trial, two people living with PSC will be members of the Trial Steering Committee.

Given the length of the trial and the nature of PSC, membership may change as some members become too sick to participate. Membership will be drawn from PSC Support's ongoing network of patients and loved ones from across the UK who are interested in being involved in research. Lay Trial Steering Committee members will join with different needs and levels of expertise, knowledge and experience, and the PPIE Lead will offer tailored training and support to members to build skills and confidence as and when required. People at risk of developing PSC cancers are a diverse population and the aim is to reflect that so that a wide range of views and perspectives are heard. To that end, focus groups will be used when necessary and collect and review optional demographic data, addressing any under-represented groups.

Allowing meeting attendance via teleconferencing widens participation as it allows people to join from all corners of the UK, including those who are clinically vulnerable to COVID. We will send printed materials to those who need them and liaise by phone call if Zoom is not accessible for some members. One lay member of the public will sit on the

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

Independent Data Monitoring Committee. They will be recruited from the People in Research and Cancer Research UK networks.

PPIE will be engaged to contribute to activities at the following milestones:

Trial Set-Up

- Review of key sections of the protocol to identify preventable barriers to recruitment and retention.
- Assist in the development of patient facing materials, such as the recruitment documents, Patient Information Sheet (PIS) and consent form to ensure ease of understanding and use of appropriate language (online focus group activity).

Ongoing Trial

- Review recruitment rate and provide feedback on the recruitment strategy as part of the Trial Steering Committee.
- Review lay interim reports (if there are any) and changes to any patient-facing documentation or communications (online focus group activity).

Trial End

- Review and assist in the development of key messages and the final lay report for dissemination to participants, the public and PSC patients including producing engaging media such as infographics and videos (online focus group activity).
- Advise on key messages and dissemination avenues (online focus group activity).

With the support of the ICTU Study Manager and Chief Investigator, the PPIE leads will prepare background pre-reading for any focus groups required and the draft the documents for review. Using online focus groups will help us to ensure a diverse range of patients' perspectives are considered throughout trial life cycle. The PPIE leads will also support the team at Investigator Meetings by providing the patient perspective on the study.

11.9.4 PPIE Evaluation and Feedback

A detailed log of PPIE activities and their impact will be kept throughout trial, to enable evaluation of PPIE impact. The log will include PPIE trial aims; PPIE methods used; PPIE results (positive/negative); extent that PPIE influenced the trial and a reflection on progress. Impact feedback to TSC and focus group members will be given throughout trial to keep engagement and motivation levels high and also after trial involvement has concluded. This feedback summary will include suggestions made; how suggestions were taken on board and overall impact on the trial.

The ICTU study manager has considerable experience working as PPIE liaison for clinical trials, supporting patients who wish to act as PPIE collaborators for trial groups and focus groups. Further support will be provided by the PPIE Lead/Chief Investigator where needed.

11.10 PUBLICATION AND DISSEMINATION POLICY

The results will be available as a lay summary (available to download or view), produced in collaboration with our PPIE members as well as links to the Open Access scientific, peer-

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

reviewed publications. Results will also be shared by the patient organisations, PSC Support and AMMF, via their newsletters, social media and Patient Information Days and conferences.

Information concerning the study, patent applications, processes, scientific data or other pertinent information is confidential and remains the property of the Sponsor. The investigator may use this information for the purposes of the study only.

It is understood by the investigator that the Sponsor will use information developed in this clinical study in connection with the development of the IMP and, therefore, may disclose it as required to other clinical investigators and to Regulatory Authorities. In order to allow the use of the information derived from this clinical study, the investigator understands that he/she has an obligation to provide complete test results and all data developed during this study to the Sponsor.

Verbal or written discussion of results prior to study completion and full reporting should only be undertaken with written consent from the Sponsor.

Therefore, all information obtained because of the study will be regarded as CONFIDENTIAL, at least until appropriate analysis and review by the investigator(s) are completed.

A Clinical Study Report summarising the study results will be prepared and submitted to the REC within a year of the end of study.

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

12 REFERENCES

1. Chapman MH, Thorburn D, Hirschfield GM, Webster GGJ, Rushbrook SM, Alexander G et al. British Society of Gastroenterology and UK-PSC guidelines for the diagnosis and management of primary sclerosing cholangitis. *Gut* 2019; Aug;68(8):1356-1378. doi: 10.1136/gutjnl-2018-317993.
2. Trivedi PJ, Crothers H, Mytton J, Bosch S, Iqbal T, Ferguson J et al. Effects of Primary Sclerosing Cholangitis on Risks of Cancer and Death in People With Inflammatory Bowel Disease, Based on Sex, Race, and Age. *Gastroenterol* 2020;159(3):915-928. doi: 10.1053/j.gastro.2020.05.049.
3. Trivedi PJ, Bowlus CL, Yimam KK, Razavi H, Estes C. et al. Epidemiology, Natural History, and Outcomes of Primary Sclerosing Cholangitis: A Systematic Review of Population-based Studies. *Clin Gastroenterol Hepatol* 2021;
4. Trivedi P & Hirschfield GM. Recent advances in clinical practice: epidemiology of autoimmune liver diseases. *Gut* 2021;70(10):1989-2003.
5. Boonstra K, Weersma RK, van Erpecum KJ, Rauws EA, Spanier BW, Poen AC et al. Population-based epidemiology, malignancy risk, and outcome of primary sclerosing cholangitis. *Hepatology* 2013 Dec;58(6):2045-55. doi: 10.1002/hep.26565.
6. Aune D, Sen A, Norat T, Riboli E, Folseraas T. Primary sclerosing cholangitis and the risk of cancer, cardiovascular disease, and all-cause mortality: a systematic review and meta-analysis of cohort studies. *Sci Rep* 2021 May 20;11(1):10646. doi: 10.1038/s41598-021-90175-w.
7. Banales JM, Marin JJG, Lamarca A, Rodrigues PM, Khan SA, Roberts LR et. al. Cholangiocarcinoma 2020: the next horizon in mechanisms and management. *Nature Rev Gastroenterol Hepatol*. 2020 Jun 30. doi: 10.1038/s41575-020-0310-z.
8. Weismuller TJ,. Patient Age, Sex, and Inflammatory Bowel Disease Phenotype Associate With Course of Primary Sclerosing Cholangitis. *Gastroenterology* 2017doi: 10.1053/j.gastro.2017.02.038.
9. Fatourou et al. OWE-4 National audit of diagnosis, management and surveillance in primary sclerosing cholangitis in the United Kingdom. *Gut* Nov 2021, 70 (Suppl 4) A9; DOI: 10.1136/gutjnl-2021-BSG.16.
10. Ali AH, Tabibian JH, Nasser-Ghods N, Lennon RJ, DeLeon T, Borad MJ et al. Surveillance for hepatobiliary cancers in patients with primary sclerosing cholangitis. *Hepatology* 2018 Jun;67(6):2338-2351. doi: 10.1002/hep.29730.
11. García Rodríguez LA, Soriano-Gabarró M, Bromley S, Lanas A, Soriano LC. New use of low-dose aspirin and risk of colorectal cancer by stage at diagnosis: a nested case-control study in UK general practice. *BMC Cancer* 2017;17(1):637. doi: 10.1186/s12885-017-3594-9.
12. Tsoi KK, Chan FC, Hirai HW, Sung JJ. Risk of gastrointestinal bleeding and benefit from colorectal cancer reduction from long-term use of low-dose aspirin: A retrospective study of 612 509 patients. *J Gastroenterol Hepatol* 2018. 33(10):1728-1736. doi: 10.1111/jgh.14261.
13. Burn J, Sheth H, Elliott F, Reed L, Macrae F, Mecklin J-P, Möslin G, McDonald FE, Bertario L et al. Cancer prevention with aspirin in hereditary colorectal cancer (Lynch syndrome), 10-year follow-up and registry-based 20-year data in the CAPP2 study: a double-blind, randomised, placebo-controlled trial. *Lancet*; 2020

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

14. NICE guideline [NG151] Colorectal cancer 2020: <https://www.nice.org.uk/guidance/ng151/>
15. Bosetti C, Santucci C, Gallus S, Martinetti M, La Vecchia C. Aspirin and the risk of colorectal and other digestive tract cancers: an updated meta-analysis through 2019. *Annals Onc* 2020; 31:5:558-568.
16. Choi J, Ghazal HM, Peeraphatdit T, Baichoo E, Addissie BD, Harmsen WS et al 2016. Aspirin use and the risk of cholangiocarcinoma; *Hepatology* 2016; 016;64(3):785-96. doi: 10.1002/hep.28529.
17. Lapumnuaypol K, Tiu A, Thongprayoon C, Wijarnpreecha K, Ungprasert P, Mao MA et al. Effects of aspirin and non-steroidal anti-inflammatory drugs on the risk of cholangiocarcinoma: a meta-analysis. *QJM*. 2019 Jun 1;112(6):421-427. doi: 10.1093/qjmed/hcz039.
18. Xiong J, Xu W, Bian J, Huang H, Bai Y, Xu Y et al. Aspirin use is associated with a reduced risk of cholangiocarcinoma: a systematic review and meta-analysis. *Cancer Manag Res*. 2018; 10: 4095–4104. doi: 10.2147/CMAR.S173197.
19. Langley RE, Burdett S, Tierney JF, Cafferty F, Parmar MKB, Venning G. Aspirin and cancer: has aspirin been overlooked as an adjuvant therapy? *Br J Cancer* 2011; 105, 1107 – 1113. doi: 10.1038/bjc.2011.289.
20. Jackson SS, Pfeiffer RM, Liu Z, Anderson LA, Tsai HT, Gadalla SM et al. Association Between Aspirin Use and Biliary Tract Cancer Survival. *JAMA Oncol*. 2019;5(12):1802-1804. doi: 10.1001/jamaoncol.2019.4328.
21. Liao S-F, Koshiol J, Huang Y-H, Jackson SS, Huang Y-H, Chan C et al. Postdiagnosis Aspirin Use Associated With Decreased Biliary Tract Cancer-Specific Mortality in a Large Nationwide Cohort. *Hepatology* 2021.
22. Simon TG, Duberg AS, Aleman S, Chung RT, Chan AT, Ludvigsson JF. Association of Aspirin with Hepatocellular Carcinoma and Liver-Related Mortality. *N Engl J Med* 2020 Mar 12;382(11):1018-1028. doi: 10.1056/NEJMoa1912035.
23. Shen X & Shen X. A potential role for aspirin in the prevention and treatment of cholangiocarcinoma. *Int J Cancer* 2021 Mar 15;148(6):1323-1330. doi: 10.1002/ijc.33323.
24. Shi T, Gong J, Fujita K, Nishiyama N, Iwama H, Liu S et al. Aspirin inhibits cholangiocarcinoma cell proliferation via cell cycle arrest in vitro and in vivo. *Int J Oncol*. 2021 Feb; 58(2): 199–210. doi: 10.3892/ijo.2020.5165.
25. Elman S, Hynan LS, Gabriel V, Mayo MJ. The 5-D itch scale: a new measure of pruritus. *Br J Dermatol*. 2010;162(3):587-93.
26. Younossi ZM, Afendy A, Stepanova M, Racila A, Nader F, Gomel R, et al. Development and validation of a primary sclerosing cholangitis-specific patient-reported outcomes instrument: The PSC PRO. *Hepatology*. 2018;68(1):155-65
27. Garza-Galvan et al. Low-dose aspirin use in the first trimester of pregnancy and odds of congenital anomalies: A meta-analysis of randomized controlled trials. *Int J Gynaecol Obstet*. 2023 Feb;160(2):526-537. doi: 10.1002/ijgo.14334. PMID: 35810407, DOI: 10.1002/ijgo.14334.
28. Roberge S, Bujold E, Nicolaides KH. Aspirin for the prevention of preterm and term preeclampsia: systematic review and metaanalysis. *Am J Obstet Gynecol* 2018;218(3):287-93 e1. doi: 10.1016/j.ajog.2017.11.56

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

29. Rolnik DL et al. Aspirin versus Placebo in Pregnancies at High Risk for Preterm Preeclampsia. N Engl J Med. 2017;377(7):613-622. doi: 10.1056/NEJMoa1704559. PMID: 28657417, DOI:10.1056/NEJMoa1704559.

13 REVISION HISTORY

Version	Date	Summary of changes
1.0	23/08/2023	N/A
2.0	20/10/2023	<ul style="list-style-type: none"> • Exclusion Criteria updated • Contraindicated section updated • Pregnancy section updated • Unblinding section updated
3.0	DRAFT	<p>General minor wording changes</p> <p>Addition of REC and ICRTN number and named Clinical Trial Monitor</p> <p>Change of Study statistician</p> <p>Change of Vendor storing scans</p> <p>Addition of vendors to take ELF test</p> <p>Addition of Trial Monitor contact details</p> <p>Minor wording changes to inclusion/exclusion criteria</p> <p>Removal of untreated thyrotoxicosis and hypothyroidism from exclusion criteria, as these are not specified in the SmPC.</p> <p>SSRI use removed as exclusion (reasons detailed in cover letter)</p> <p>Trial procedure changes: stool samples no longer sent to GP; standard of care blood tests clarified as every 6 months.</p> <p>Stratification will be done by ELF test or fibroscan, if ELF test is not available</p> <p>1Stratification category for PSC duration changed</p> <p>6-month visit can only be performed with a +5 day window and not a +14 day window</p>

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

		<p>Clarification how to give the one month questionnaire</p> <p>Contraindication updates: added CTIMP participation and use of aspirin; removed NSAID use and SSRI</p> <p>Clarification on withdrawal procedures</p> <p>Removal of Annual Progress Reports as this is no longer an HRA requirement.</p> <p>Clarification that only a doctor can take consent</p>
--	--	--

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

14 SIGNATURE PAGE 1 (CHIEF INVESTIGATORS)

The signature below constitutes approval of this protocol by the signatory, on behalf of the Protocol Development Group, and provides the necessary assurances that this study will be conducted according to all stipulations of the protocol including all statements regarding confidentiality.

Study Title: Asp-PSC Trial: Effect of Aspirin on Reducing Cancer & Improving Outcomes in Primary Sclerosing Cholangitis

Protocol Number: C/45/2022

Signed:

DocuSigned by:

shahid khan

059CEF8AD8D949A...

Professor Shahid Khan

Date: 13-May-2025

Signed:

DocuSigned by:

Simon Rushbrook

AFBC959C5D9C4E4...

Dr Simon Rushbrook

Date: 16-May-2025

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------


15 SIGNATURE PAGE 2 (SPONSOR)

The signature below constitute approval of this protocol by the signatory.

Study Title: Asp-PSC Trial: Effect of Aspirin on Reducing Cancer & Improving Outcomes in Primary Sclerosing Cholangitis

Protocol Number: C/45/2022

Signed:

Signed by:

385B924F3D2F40A...

Imperial College London
Research Governance and Integrity team
Room 221, Medical School Building
St Marys Campus, Norfolk Place
London W2 1PG
United Kingdom

Date:

13-May-2025

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

16 SIGNATURE PAGE 3 (STATISTICIAN)

The signature below constitute approval of this protocol by the signatory.

Study Title: Asp-PSC Trial: Effect of Aspirin on Reducing Cancer & Improving Outcomes in Primary Sclerosing Cholangitis

Protocol Number: C/45/2022

Signed: _____

Mr Nicholas Johnson
Imperial College London

Date: _____

Rachel
Phillip
S

Digitally signed
by Rachel
Phillips
Date:
2025.05.13
08:52:58 [®]
+01'00'

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

17 SIGNATURE PAGE 4 (PRINCIPAL INVESTIGATOR)

The signature of the below constitutes agreement of this protocol by the signatory and provides the necessary assurance that this study will be conducted at his/her investigational site according to all stipulations of the protocol including all statements regarding confidentiality.

Study Title: Asp-PSC Trial: Effect of Aspirin on Reducing Cancer & Improving Outcomes in Primary Sclerosing Cholangitis

Protocol Number: C/45/2022

Address of Institution: _____

Signed: _____

Print Name and Title: _____

Date: _____

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

18 APPENDICES

Appendix 1: PSC-PRO (26)

The PSC-PRO is a validates disease specific questionnaire frequently employed in clinical trials of patients with PSC to derive quality of life outcomes. It covers 29 items across two main outcomes: 'symptoms' and 'impact of symptoms'.

Module 1 – PSC Symptoms

Scores: 0 = “no symptoms” to 10 = “symptoms as bad as you could imagine”; recall period 24 hours.

1. Are you currently experiencing a flare-up of your PSC symptoms (also known as acute cholangitis or an infection in the bile ducts)? Y/N
2. Over the past 24 hours, how bad was the pain in your upper abdomen due to PSC at its worst?
3. Over the past 24 hours, how bad was the discomfort in your upper abdomen due to PSC at its worst?
4. Over the past 24 hours, how bad was your itching at its worst?
5. Over the past 24 hours, how bad was your physical tiredness at its worst?
6. Over the past 24 hours, how bad was the yellowing of your eyes or skin at its worst?
7. Over the past 24 hours, how bad was your difficulty concentrating at its worst?
8. Over the past 24 hours, how bad was your nausea at its worst?
9. Over the past 24 hours, how bad was your mental tiredness at its worst?
10. Over the past 24 hours, how bad was the darkening (brown or tea color) of your urine at its worst?
11. Over the past 24 hours, how bad were your fever (high temperature) symptoms?
12. Over the past 24 hours, how bad were your chills?
13. Over the past 24 hours, how bad were your sweats?

Module 2 - Impacts of Symptoms

Scores: 1 = “Never” to 5 = “Always”; recall period 7 days.

Physical Function

1. Over the past 7 days, how often was it difficult for you to go up or down stairs because of your PSC?
2. Over the past 7 days, how often did you have to rest or take breaks because of your PSC?
3. Over the past 7 days, how often were you limited in what you could do physically because of your PSC?
4. Over the past 7 days, how often was it difficult for you to lift or carry objects because of your PSC?

Activities of Daily Living

5. Over the past 7 days, how often was it difficult for you to complete household chores because of your PSC?
6. Over the past 7 days, how often did you have difficulty falling or staying asleep because of your PSC?
7. Over the past 7 days, how often was it difficult to run errands or get things done outside the home because of your PSC?
8. Over the past 7 days, how often did you avoid certain foods because of your PSC?

Work Productivity

9. Are you currently employed or self-employed? Y/N (if N then skip the section)
10. Over the past 7 days, how often did PSC interfere with your productivity at work?
11. Over the past 7 days, how often did you limit your hours at work because of your PSC?
12. Over the past 7 days, how often was it an extra effort to do your work because of your PSC?
13. Over the past 7 days, how often was it difficult to concentrate on your work because of your PSC?

Role Function

14. Over the past 7 days, how often did you need to ask for help around your home because of your PSC?
15. Over the past 7 days, how often did you feel like a burden on your family because of your PSC?
16. Over the past 7 days, how often were you limited in the activities you could do with your family because of your PSC?
17. Over the past 7 days, how often was it difficult to enjoy time with your family because of your PSC?

Emotional Impact

6

18. Over the past 7 days, how often were you worried about your health because of your PSC?
19. Over the past 7 days, how often were you depressed about your health because of your PSC?
20. Over the past 7 days, how often did you feel emotional stress because of your PSC?

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

21. Over the past 7 days, how often did you feel scared about the future because of your PSC?

Social/Leisure Impact

22. Over the past 7 days, how often did you limit your social activities because of your PSC?

23. Over the past 7 days, how often did you limit the things you did for enjoyment because of your PSC?

24. Over the past 7 days, how often did you isolate yourself from other people because of your PSC?

25. Over the past 7 days, how often did you have limited energy for sexual activity because of your PSC?

Quality of Life

26. Over the past 7 days, how often was it difficult to work towards your goals in life because of your PSC?

27. Over the past 7 days, how often were you unable to enjoy life because of your PSC?

28. Over the past 7 days, how often did you feel like you missed out on meaningful activities because of your PSC?

29. Over the past 7 days, how often did you feel discouraged about your quality of life because of your PSC?

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

Appendix 2: SF-36

SF-36 QUESTIONNAIRE

Name: _____ Ref. Dr: _____ Date: _____
ID#: _____ Age: _____ Gender: M / F

Please answer the 36 questions of the **Health Survey** completely, honestly, and without interruptions.

GENERAL HEALTH:

In general, would you say your health is:

Excellent Very Good Good Fair Poor

Compared to one year ago, how would you rate your health in general now?

Much better now than one year ago

Somewhat better now than one year ago

About the same

Somewhat worse now than one year ago

Much worse than one year ago

LIMITATIONS OF ACTIVITIES:

The following items are about activities you might do during a typical day. Does your health now limit you in these activities? If so, how much?

Vigorous activities, such as running, lifting heavy objects, participating in strenuous sports.

Yes, Limited a lot Yes, Limited a Little No, Not Limited at all

Moderate activities, such as moving a table, pushing a vacuum cleaner, bowling, or playing golf

Yes, Limited a Lot Yes, Limited a Little No, Not Limited at all

Lifting or carrying groceries

Yes, Limited a Lot Yes, Limited a Little No, Not Limited at all

Climbing several flights of stairs

Yes, Limited a Lot Yes, Limited a Little No, Not Limited at all

Climbing one flight of stairs

Yes, Limited a Lot Yes, Limited a Little No, Not Limited at all

Bending, kneeling, or stooping

Yes, Limited a Lot Yes, Limited a Little No, Not Limited at all

Walking more than a mile

Yes, Limited a Lot Yes, Limited a Little No, Not Limited at all

Walking several blocks

Yes, Limited a Lot Yes, Limited a Little No, Not Limited at all

Walking one block

Yes, Limited a Lot Yes, Limited a Little No, Not Limited at all

Bathing or dressing yourself

Yes, Limited a Lot Yes, Limited a Little No, Not Limited at all

PHYSICAL HEALTH PROBLEMS:

During the past 4 weeks, have you had any of the following problems with your work or other regular daily activities as a result of your physical health?

Cut down the amount of time you spent on work or other activities

Yes No

Accomplished less than you would like

Yes No

Were limited in the kind of work or other activities

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

Yes No

Had difficulty performing the work or other activities (for example, it took extra effort)

Yes No

EMOTIONAL HEALTH PROBLEMS:

During the past 4 weeks, have you had any of the following problems with your work or other regular daily activities as

a result of any emotional problems (such as feeling depressed or anxious)?

Cut down the amount of time you spent on work or other activities

Yes No

Accomplished less than you would like

Yes No

Didn't do work or other activities as carefully as usual

Yes No

SOCIAL ACTIVITIES:

Emotional problems interfered with your normal social activities with family, friends, neighbors, or groups?

Not at all Slightly Moderately Severe Very Severe

PAIN:

How much bodily pain have you had during the past 4 weeks?

None Very Mild Mild Moderate Severe Very Severe

During the past 4 weeks, how much did pain interfere with your normal work (including both work outside the

home and housework)?

Not at all A little bit Moderately Quite a bit Extremely

ENERGY AND EMOTIONS:

These questions are about how you feel and how things have been with you during the last 4 weeks.

For each

question, please give the answer that comes closest to the way you have been feeling.

Did you feel full of pep?

All of the time

Most of the time

A good Bit of the Time

Some of the time

A little bit of the time

None of the Time

Have you been a very nervous person?

All of the time

Most of the time

A good Bit of the Time

Some of the time

A little bit of the time

None of the Time

Have you felt so down in the dumps that nothing could cheer you up?

All of the time

Most of the time

A good Bit of the Time

Some of the time

A little bit of the time

None of the Time

Have you felt calm and peaceful?

Confidential

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

All of the time
Most of the time
A good Bit of the Time
Some of the time
A little bit of the time
None of the Time

Did you have a lot of energy?

All of the time
Most of the time
A good Bit of the Time
Some of the time
A little bit of the time
None of the Time

Have you felt downhearted and blue?

All of the time
Most of the time
A good Bit of the Time
Some of the time
A little bit of the time
None of the Time

Did you feel worn out?

All of the time
Most of the time
A good Bit of the Time
Some of the time
A little bit of the time
None of the Time

Have you been a happy person?

All of the time
Most of the time
A good Bit of the Time
Some of the time
A little bit of the time
None of the Time

Did you feel tired?

All of the time
Most of the time
A good Bit of the Time
Some of the time
A little bit of the time
None of the Time

SOCIAL ACTIVITIES:

During the past 4 weeks, how much of the time has your physical health or emotional problems interfered with

your social activities (like visiting with friends, relatives, etc.)?

All of the time
Most of the time
Some of the time
A little bit of the time
None of the Time

GENERAL HEALTH:

How true or false is each of the following statements for you?

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

I seem to get sick a little easier than other people
Definitely true Mostly true Don't know Mostly false Definitely false

I am as healthy as anybody I know
Definitely true Mostly true Don't know Mostly false Definitely false

I expect my health to get worse
Definitely true Mostly true Don't know Mostly false Definitely false

My health is excellent
Definitely true Mostly true Don't know Mostly false Definitely false

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

Appendix 3 5-D Pruritus scale (25):

5-D Pruritus Scale

1. **Duration:** During the last 2 weeks, how many hours a day have you been itching?

Less than 6hrs/day	6-12 hrs/day	12-18 hrs/day	18-23 hrs/day	All day
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
1	2	3	4	5

2. **Degree:** Please rate the intensity of your itching over the past 2 weeks

Not present	Mild	Moderate	Severe	Unbearable
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
1	2	3	4	5

3. **Direction:** Over the past 2 weeks has your itching gotten better or worse compared to the previous month?

Completely resolved	Much better, but still present	Little bit better, but still present	Unchanged	Getting worse
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
1	2	3	4	5

4. **Disability:** Rate the impact of your itching on the following activities over the last 2 weeks

	Never affects sleep	Occasionally delays falling asleep	Frequently delays falling asleep	Delays falling asleep and occasionally wakes me up at night	Delays falling asleep and frequently wakes me up at night
Sleep	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
	1	2	3	4	5
	N/A	Never affects this activity	Rarely affects this activity	Occasionally affects this activity	Frequently affects this activity
Leisure/Social	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
		1	2	3	4
Housework/Errands	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
		1	2	3	4
Work/School	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
		1	2	3	4

5. **Distribution:** Mark whether itching has been present in the following parts of your body over the last 2 weeks. If a body part is not listed, choose the one that is closest anatomically.

Head/Scalp	<input type="checkbox"/>	Soles	<input type="checkbox"/>
Face	<input type="checkbox"/>	Palms	<input type="checkbox"/>
Chest	<input type="checkbox"/>	Tops of Hands/Fingers	<input type="checkbox"/>
Abdomen	<input type="checkbox"/>	Forearms	<input type="checkbox"/>
Back	<input type="checkbox"/>	Upper Arms	<input type="checkbox"/>
Buttocks	<input type="checkbox"/>	Points of Contact w/ Clothing (e.g waistband, undergarment)	<input type="checkbox"/>
Thighs	<input type="checkbox"/>	Groin	<input type="checkbox"/>
Lower legs	<input type="checkbox"/>		
Tops of Feet/Toes	<input type="checkbox"/>		

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

Appendix 4: Short Inflammatory Bowel Disease (SIBDQ)

Patient Name _____ Date _____

This questionnaire is designed to find out how you have been feeling during the last 2 weeks. You will be asked about symptoms you have been having as a result of your inflammatory bowel disease, the way you have been feeling in general, and how your mood has been. Please check the box of your choice below each question.

1. How often has the feeling of fatigue or being tired and worn out been a problem for you during the past 2 weeks?

- ☐ All of the time
- ☐ Most of the time
- ☐ A good bit of the time
- ☐ Some of the time
- ☐ A little of the time
- ☐ Hardly any of the time
- ☐ None of the time

2. How often during the last 2 weeks have you delayed or canceled a social engagement because of your bowel problem?

- ☐ All of the time
- ☐ Most of the time
- ☐ A good bit of the time
- ☐ Some of the time
- ☐ A little of the time
- ☐ Hardly any of the time
- ☐ None of the time

3. As a result of your bowel problems, how much difficulty did you experience doing leisure or sports activities during the past 2 weeks?

- ☐ A great deal of difficulty; activities made impossible
- ☐ A lot of difficulty
- ☐ A fair bit of difficulty
- ☐ Some difficulty
- ☐ A little difficulty
- ☐ Hardly any difficulty
- ☐ No difficulty; the bowel problem did not limit sports or leisure activities

4. How often during the past 2 weeks have you been troubled by pain in the abdomen?

- ☐ All of the time
- ☐ Most of the time
- ☐ A good bit of the time
- ☐ Some of the time
- ☐ A little of the time
- ☐ Hardly any of the time
- ☐ None of the time

5. How often during the past 2 weeks have you felt depressed or discouraged?

- ☐ All of the time
- ☐ Most of the time
- ☐ A good bit of the time
- ☐ Some of the time

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

- ☐ A little of the time
- ☐ Hardly any of the time
- ☐ None of the time

6. Overall, in the past 2 weeks, how much of a problem have you had with passing large amounts of gas?

- ☐ A major problem
- ☐ A big problem
- ☐ A significant problem
- ☐ Some problem
- ☐ A little trouble
- ☐ Hardly any trouble
- ☐ No trouble

7. Overall, in the past 2 weeks, how much of a problem have you had maintaining or getting to the weight you would like to be?

- ☐ A major problem
- ☐ A big problem
- ☐ A significant problem
- ☐ Some problem
- ☐ A little trouble
- ☐ Hardly any trouble
- ☐ No trouble

8. How often during the past 2 weeks have you felt relaxed and free of tension?

- ☐ All of the time
- ☐ Most of the time
- ☐ A good bit of the time
- ☐ Some of the time
- ☐ A little of the time
- ☐ Hardly any of the time
- ☐ None of the time

9. How much of the time during the past 2 weeks have you been troubled by a feeling of having to go to the bathroom even though your bowels were empty?

- ☐ All of the time
- ☐ Most of the time
- ☐ A good bit of the time
- ☐ Some of the time
- ☐ A little of the time
- ☐ Hardly any of the time
- ☐ None of the time

10. How often during the past 2 weeks have you felt angry as a result of your bowel problem?

- ☐ All of the time
- ☐ Most of the time
- ☐ A good bit of the time
- ☐ Some of the time
- ☐ A little of the time
- ☐ Hardly any of the time
- ☐ None of the time

Version 2

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

Irvine, E. J., et. al. The short inflammatory bowel disease questionnaire: A quality of life instrument for community physicians managing inflammatory bowel disease. Am J Gastroenterology. 1996; 91(8):1571-8 Aug.

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

Appendix 5: CLDQ-PSC

Supplementary Table 2. The list of questions included in the final version of the CLDQ-PSC instrument.

1. How much of the time during the last two weeks have you been troubled by a feeling of abdominal bloating?
2. How much of the time have you been tired or fatigued during the last two weeks?
3. **How much of the time during the last 2 weeks have you experienced bodily pain?**
4. How often during the last two weeks have you felt sleepy during the day?
5. How much of the time during the last two weeks have you experienced abdominal pain?
6. How much of the time in the last two weeks have you been bothered by having decreased strength?
7. How often during the last two weeks have you felt anxious?
8. How often during the last 2 weeks have you felt a decreased level of energy?
9. How much of the time during the last two weeks have you felt unhappy?
10. How often during the last two weeks have you felt drowsy?
11. How often during the last two weeks have you been irritable?
12. How much of the time during the last two weeks have you had difficulty sleeping at night?
13. How much of the time during the last two weeks have you been troubled by a feeling of abdominal discomfort?
14. How much of the time during the last two weeks have you been worried about the impact your liver disease has on your family?
15. How much of the time during the last two weeks have you had mood swings?
16. How much of the time during the last two weeks have you been unable to fall asleep at night?
17. How often during the last two weeks have you had muscle cramps?
18. How much of the time during the last two weeks have you been worried that your symptoms will develop into major problems?
19. How much of the time during the last two weeks have you felt depressed?
20. How much of the time during the last two weeks have you been worried about your condition getting worse?
21. How much of the time during the last two weeks have you had problems concentrating?
22. How much of the time have you been troubled by itching during the last two weeks?
23. How much of the time during the last two weeks have you been worried about never feeling any better?
24. How much of the time during the last two weeks have you been concerned about the availability of a liver if you need a liver transplant?

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

Appendix 6: PRO-2

Please indicate how you perceive your stool frequency (Based on the last 3 days)
<input type="checkbox"/> Normal
<input type="checkbox"/> 1–2 more stools than normal
<input type="checkbox"/> 3–4 more stools than normal
<input type="checkbox"/> 5 + more stools than normal
Please indicate the severity of your rectal bleeding (Based on the last 3 days)
<input type="checkbox"/> No blood seen
<input type="checkbox"/> Streaks of blood seen with stools for half of the time
<input type="checkbox"/> Obvious blood with stool most of the time
<input type="checkbox"/> Blood alone passed (with no stool)

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

Appendix 7:COMMON TERMINOLOGY CRITERIA FOR ADVERSE EVENTS Version 5.0 (CTCAE)

Toxicities will be recorded according to CTCAE version 5.0. The full documents is available from the National Cancer Institute (NCI) website and the quick reference can be viewed via the following link:

https://ctep.cancer.gov/protocoldevelopment/electronic_applications/docs/CTCAE_v5_Quick_Reference_5x7.pdf

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

Appendix 8: TOKYO GUIDELINES 2018 DEFINITION OF ACUTE CHOLANGITIS³¹

Diagnosis is made on account of a combination of clinical, biochemical and radiological parameters as established on the basis of the consensus reached in the International Consensus Meeting held in Tokyo 2018.

Evidence of	Parameters	Thresholds
A. Systemic Inflammation	<ul style="list-style-type: none"> Fever/Rigors Biochemic evidence of inflammatory response 	<ul style="list-style-type: none"> Temperature >38°C Leucopenia (<4 x1000 µl), leucocytosis (>10 x 1000µl) or elevated CRP (>1mg/dl)
B. Cholestasis	<ul style="list-style-type: none"> Jaundice Abnormal liver function tests on biochemistry 	<ul style="list-style-type: none"> Elevated bilirubin (>2g/dl) Elevated ALP/AST/ALT/GGT > 1.5x ULN
C. Imaging (Ultrasound, endoscopic ultrasound, MRCP)	<ul style="list-style-type: none"> Dilated biliary system Organic pathology seen on imaging 	<ul style="list-style-type: none"> Bile duct >7mm Evidence of Stone, stricture, or obstruction

Tokyo Guidelines score Interpretation:

Definite diagnosis	one item in A, one item in B and one item in C
Suspected diagnosis	one item in A + one item in either B or C

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

Appendix 9: Bristol Laboratories Ltd SmPC Version 10, January 2021

Side effects are grouped on the basis of System Organ Class. Within each system organ class the frequencies are defined as: very common ($\geq 1/10$), common ($\geq 1/100$ to $< 1/10$), uncommon ($\geq 1/1,000$ to $< 1/100$), rare ($\geq 1/10,000$ to $< 1/1,000$), very rare ($< 1/10,000$) and not known (cannot be estimated from the available data).

Blood and lymphatic system disorders

Common:

Increased bleeding tendencies.

Rare:

Thrombocytopenia, granulocytosis, aplastic anaemia.

Not known:

Cases of bleeding with prolonged bleeding time such as epistaxis and gingival bleeding.

Symptoms may persist for a period of 4–8 days after acetylsalicylic acid discontinuation. As a result there may be an increased risk of bleeding during surgical procedures.

Existing (haematemesis, melaena) or occult gastrointestinal bleeding, which may lead to iron deficiency anaemia (more common at higher doses).

Immune system disorders

Rare:

Hypersensitivity reactions, angio-oedema, allergic oedema, anaphylactic reactions including shock.

Not known:

Hyperuricemia.

Metabolism and digestive system disorders

Nervous system disorders

Rare:

Intracranial haemorrhage

Not known:

Headache, vertigo

Not known:

Reduced hearing ability; tinnitus

Rare:

Hemorrhagic vasculitis.

Uncommon:

Rhinitis, dyspnoea.

Rare:

Bronchospasm, asthma attacks.

Rare: Menorrhagia

Ear and labyrinth disorders

Vascular disorders

Respiratory, thoracic and mediastinal disorders

Reproductive

System and mammary disorders

Gastrointestinal disorders

Common:

Dyspepsia.

Rare:

Severe gastrointestinal haemorrhage, nausea, vomiting

Not known:

Gastric or duodenal ulcers and perforation, diarrhoea

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

Hepatobiliary disorders

*Not known:
Hepatic insufficiency*

Skin and subcutaneous tissue disorders

*Uncommon:
Urticaria.
Rare:
Steven-Johnsons syndrome, Lyells syndrome, purpura, erythema nodosum, erythema multiforme.*

Renal and urinary tract disorders

Not known: Impaired renal function, salt and water retention.

Drug interactions

Interaction with Other Medicinal Products and Other Forms of Interaction

Contraindicated combinations

Methotrexate (used at doses >15 mg/week):

The combined drugs, methotrexate and acetylsalicylic acid, enhance haematological toxicity of methotrexate due to the decreased renal clearance of methotrexate by acetylsalicylic acid. Therefore, the concomitant use of methotrexate (at doses >15 mg/week) with Aspirin 75 mg tablets is contraindicated (

Not recommended combinations:

Uricosuric agents, e.g. probenecid and sulfinpyrazone

Salicylates reverse the effect of probenecid and sulfinpyrazone. The combination should be avoided

Combinations requiring precautions for use or to be taken into account:

Anticoagulants e.g. coumarin, heparin, warfarin and phenindione:

Increased risk of bleeding due to inhibited thrombocyte function, injury of the duodenal mucosa and displacement of oral anticoagulants from their plasma protein binding sites. The bleeding time should be monitored

Anti-platelet agents (e.g. clopidogrel and dipyridamole) and selective serotonin re-uptake inhibitors (SSRIs; such as sertraline or paroxetine):

Increased risk of gastrointestinal bleeding

Salicylics may increase the hypoglycaemic effect of sulphonylureas.

Digoxin and lithium:

Acetylsalicylic acid impairs the renal excretion of digoxin and lithium, resulting in increased plasma concentrations. Monitoring of plasma concentrations of digoxin and lithium is recommended when initiating and terminating treatment with acetylsalicylic acid. Dose adjustment may be necessary

Diuretics and antihypertensives:

NSAIDs may decrease the antihypertensive effects of diuretics and other antihypertensive agents. As for other NSAIDs concomitant administration with ACE-inhibitors increases the risk of acute renal insufficiency.

Diuretics: Risk of acute renal failure due to the decreased glomerular filtration via decreased renal prostaglandin synthesis. Hydrating the patient and monitoring renal function at the start of the treatment is recommended.

Carbonic anhydrase inhibitors (acetazolamide):

May result in severe acidosis and increased central nervous system toxicity.

Imperial Clinical Trials Unit	Asp-PSC Protocol Number: C/45/2022	V3.0 18.02.2025
-------------------------------	---------------------------------------	-----------------

Systemic Corticosteroids:

The risk of gastrointestinal bleeding and ulceration is increased when acetylsalicylic acid and corticosteroids are co-administered

Methotrexate (used at doses <15 mg/week):

The combined drugs, methotrexate and acetylsalicylic acid, may increase haematological toxicity of methotrexate due to decreased renal clearance of methotrexate by acetylsalicylic acid. Weekly blood count checks should be done during the first weeks of the combination. Enhanced monitoring should take place in the presence of even mildly impaired renal function, as well, as in elderly.

Other non-steroidal anti-inflammatory drugs (NSAIDs):

Increased risk of ulcerations and gastrointestinal bleeding due to synergistic effects.

Ibuprofen:

Experimental data suggest that ibuprofen may inhibit the effect of low dose aspirin on platelet aggregation when they are dosed concomitantly. However, the limitations of these data and the uncertainties regarding extrapolation of ex vivo data to the clinical situation imply that no firm conclusions can be made for regular ibuprofen use, and no clinically relevant effect is considered to be likely for occasional ibuprofen use

Ciclosporin, tacrolimus:

Concomitant use of NSAIDs and ciclosporin or tacrolimus may increase the nephrotoxic effect of ciclosporin and tacrolimus. The renal function should be monitored in case of concomitant use of these agents and acetylsalicylic acid.

Valproate

Acetylsalicylic acid has been reported to decrease the binding of valproate to serum albumin, thereby increasing its free plasma concentrations at steady state.

Phenytoin (Anti-epileptics):

Salicylate diminishes the binding of phenytoin to plasma albumin. This may lead to decreased total phenytoin levels in plasma, but increased free phenytoin fraction. The unbound concentration, and thereby the therapeutic effect, does not appear to be significantly altered.

Alcohol:

Concomitant administration of alcohol and acetylsalicylic acid increases the risk of gastrointestinal bleeding.

Metamizole:

Metamizole may reduce the effect of acetylsalicylic acid on platelet aggregation, when taken concomitantly. Therefore, this combination should be used with caution in patients taking low dose aspirin for cardioprotection.

Certificate Of Completion

Envelope Id: F2C3110F-9A81-438D-A827-47417883E069

Status: Completed

Subject: Complete with Docusign: Asp-PSC_ Protocol_v3.0_18.02.25 Clean.pdf

Source Envelope:

Document Pages: 88

Signatures: 3

Envelope Originator:

Certificate Pages: 5

Initials: 0

Menachi Reddi

AutoNav: Enabled

m.reddi@imperial.ac.uk

Envelopeld Stamping: Enabled

IP Address: 2a0c:5bc0:40:2e

Time Zone: (UTC) Dublin, Edinburgh, Lisbon, London

Record Tracking

Status: Original

Holder: Menachi Reddi

Location: DocuSign

13-May-2025 | 11:32

m.reddi@imperial.ac.uk

Signer Events

Nicholson, Ruth M

r.nicholson@imperial.ac.uk

Security Level: Email, Account Authentication
(None)

Signature

Signed by:

Nicholson, Ruth M
385B924F3D2F40A...

Timestamp

Sent: 13-May-2025 | 11:43

Viewed: 13-May-2025 | 11:45

Signed: 13-May-2025 | 11:45

Signature Adoption: Pre-selected Style

Using IP Address: 31.54.38.216

Electronic Record and Signature Disclosure:

Not Offered via Docusign

shahid khan

shahid.khan1@nhs.net

Consultant and CI

Security Level: Email, Account Authentication
(None)

DocuSigned by:

shahid khan
059CEF8AD8D949A...

Sent: 13-May-2025 | 11:43

Viewed: 13-May-2025 | 11:51

Signed: 13-May-2025 | 11:51

Signature Adoption: Pre-selected Style

Using IP Address: 129.31.6.66

Electronic Record and Signature Disclosure:

Accepted: 13-May-2025 | 11:51

ID: 9497158e-a2e2-41e2-a70b-f7656eaeae78

Simon Rushbrook

SIMON.RUSHBROOK@nhs.uk

PI

Security Level: Email, Account Authentication
(None)

DocuSigned by:

Simon Rushbrook
AFBC959C5D9C4E4...

Sent: 13-May-2025 | 11:43

Resent: 16-May-2025 | 15:08

Viewed: 16-May-2025 | 15:15

Signed: 16-May-2025 | 15:16

Signature Adoption: Pre-selected Style

Using IP Address: 208.127.198.60

Electronic Record and Signature Disclosure:

Accepted: 11-Aug-2023 | 11:01

ID: 68696318-1101-4e62-8386-7d9167ab6c38

In Person Signer Events

Signature

Timestamp

Editor Delivery Events

Status

Timestamp

Agent Delivery Events

Status

Timestamp

Intermediary Delivery Events

Status

Timestamp

Certified Delivery Events

Status

Timestamp

Carbon Copy Events

Status

Timestamp

Witness Events	Signature	Timestamp
----------------	-----------	-----------

Notary Events	Signature	Timestamp
---------------	-----------	-----------

Envelope Summary Events	Status	Timestamps
-------------------------	--------	------------

Envelope Sent	Hashed/Encrypted	13-May-2025 11:44
Certified Delivered	Security Checked	16-May-2025 15:15
Signing Complete	Security Checked	16-May-2025 15:16
Completed	Security Checked	16-May-2025 15:16

Payment Events	Status	Timestamps
----------------	--------	------------

Electronic Record and Signature Disclosure
--

ELECTRONIC RECORD AND SIGNATURE DISCLOSURE

From time to time, Imperial College London (we, us or Company) may be required by law to provide to you certain written notices or disclosures. Described below are the terms and conditions for providing to you such notices and disclosures electronically through the DocuSign system. Please read the information below carefully and thoroughly, and if you can access this information electronically to your satisfaction and agree to this Electronic Record and Signature Disclosure (ERSD), please confirm your agreement by selecting the check-box next to 'I agree to use electronic records and signatures' before clicking 'CONTINUE' within the DocuSign system.

Getting paper copies

At any time, you may request from us a paper copy of any record provided or made available electronically to you by us. You will have the ability to download and print documents we send to you through the DocuSign system during and immediately after the signing session and, if you elect to create a DocuSign account, you may access the documents for a limited period of time (usually 30 days) after such documents are first sent to you. After such time, if you wish for us to send you paper copies of any such documents from our office to you, you will be charged a \$0.00 per-page fee. You may request delivery of such paper copies from us by following the procedure described below.

Withdrawing your consent

If you decide to receive notices and disclosures from us electronically, you may at any time change your mind and tell us that thereafter you want to receive required notices and disclosures only in paper format. How you must inform us of your decision to receive future notices and disclosure in paper format and withdraw your consent to receive notices and disclosures electronically is described below.

Consequences of changing your mind

If you elect to receive required notices and disclosures only in paper format, it will slow the speed at which we can complete certain steps in transactions with you and delivering services to you because we will need first to send the required notices or disclosures to you in paper format, and then wait until we receive back from you your acknowledgment of your receipt of such paper notices or disclosures. Further, you will no longer be able to use the DocuSign system to receive required notices and consents electronically from us or to sign electronically documents from us.

All notices and disclosures will be sent to you electronically

Unless you tell us otherwise in accordance with the procedures described herein, we will provide electronically to you through the DocuSign system all required notices, disclosures, authorizations, acknowledgements, and other documents that are required to be provided or made available to you during the course of our relationship with you. To reduce the chance of you inadvertently not receiving any notice or disclosure, we prefer to provide all of the required notices and disclosures to you by the same method and to the same address that you have given us. Thus, you can receive all the disclosures and notices electronically or in paper format through the paper mail delivery system. If you do not agree with this process, please let us know as described below. Please also see the paragraph immediately above that describes the consequences of your electing not to receive delivery of the notices and disclosures electronically from us.

How to contact Imperial College London:

You may contact us to let us know of your changes as to how we may contact you electronically, to request paper copies of certain information from us, and to withdraw your prior consent to receive notices and disclosures electronically as follows:

To contact us by email send messages to: andrew.mclean@imperial.ac.uk

To advise Imperial College London of your new email address

To let us know of a change in your email address where we should send notices and disclosures electronically to you, you must send an email message to us at andrew.mclean@imperial.ac.uk and in the body of such request you must state: your previous email address, your new email address. We do not require any other information from you to change your email address.

If you created a DocuSign account, you may update it with your new email address through your account preferences.

To request paper copies from Imperial College London

To request delivery from us of paper copies of the notices and disclosures previously provided by us to you electronically, you must send us an email to andrew.mclean@imperial.ac.uk and in the body of such request you must state your email address, full name, mailing address, and telephone number. We will bill you for any fees at that time, if any.

To withdraw your consent with Imperial College London

To inform us that you no longer wish to receive future notices and disclosures in electronic format you may:

- i. decline to sign a document from within your signing session, and on the subsequent page, select the check-box indicating you wish to withdraw your consent, or you may;
- ii. send us an email to andrew.mclean@imperial.ac.uk and in the body of such request you must state your email, full name, mailing address, and telephone number. We do not need any other information from you to withdraw consent.. The consequences of your withdrawing consent for online documents will be that transactions may take a longer time to process..

Required hardware and software

The minimum system requirements for using the DocuSign system may change over time. The current system requirements are found here: <https://support.docusign.com/guides/signer-guide-signing-system-requirements>.

Acknowledging your access and consent to receive and sign documents electronically

To confirm to us that you can access this information electronically, which will be similar to other electronic notices and disclosures that we will provide to you, please confirm that you have read this ERSD, and (i) that you are able to print on paper or electronically save this ERSD for your future reference and access; or (ii) that you are able to email this ERSD to an email address where you will be able to print on paper or save it for your future reference and access. Further, if you consent to receiving notices and disclosures exclusively in electronic format as described herein, then select the check-box next to 'I agree to use electronic records and signatures' before clicking 'CONTINUE' within the DocuSign system.

By selecting the check-box next to 'I agree to use electronic records and signatures', you confirm that:

- You can access and read this Electronic Record and Signature Disclosure; and
- You can print on paper this Electronic Record and Signature Disclosure, or save or send this Electronic Record and Disclosure to a location where you can print it, for future reference and access; and
- Until or unless you notify Imperial College London as described above, you consent to receive exclusively through electronic means all notices, disclosures, authorizations, acknowledgements, and other documents that are required to be provided or made available to you by Imperial College London during the course of your relationship with Imperial College London.